

**M1.(a)** any **one** from:

- there was a flame
  - energy was given out
  - a new substance was formed
  - the magnesium turned into a (white) powder
- answers must be from the figure*

1

(b) Magnesium oxide

1

(c) The reaction has a high activation energy

1

(d) 9

1

(e) They have a high surface area to volume ratio

1

(f) any **one** from:

- Better coverage
- More protection from the Sun's ultraviolet rays

1

(g) any **one** from:

- Potential cell damage to the body
- Harmful effects on the environment

1

(h) indication of  $\frac{1}{1.6} = 0.625$

**and**

use of indices  $10^{-9} - 10^{-6} = 10^3$

*Both steps must be seen to score first mark*

**1**

$0.625 \times 1000 = 625$  (times bigger)

**1**

**[9]**

M2.(a) (i) 11

1

(ii) 4620 (J)

*correct answer gains 2 marks with or without working*

*allow 4.62kJ for 2 marks*

*if answer is incorrect:*

*100 × 4.2 × 11 gains 1 mark*

**or**

*100 × 4.2 × (their temp. rise) gains 1 mark*

**or**

*100 × 4.2 × (their temp. rise) correctly calculated gains 2 marks*

2

(b) the temperature increases

*allow gets hotter*

*allow heat / energy is given off*

1

(c) (i) (energy of) products lower than (energy of) reactants

*allow converse*

*allow arrow C points downwards*

1

(ii) A

1

[6]

**M3.(a)** heat / energy

1

given out / transfers to surroundings

*the mark for given out / transfers to cannot be awarded without  
heat / energy*

*allow given off*

1

(b) (i) decreases

1

increases

1

(ii) it gives the particles more energy

1

it makes the particles move faster

1

[6]

M4.	(a)	22	1
	(b)	(i) exothermic	1
		(ii) C	1
		gives out most heat energy <i>accept has largest temperature change / increase</i> <i>allow has highest (final) temperature or hottest</i>	1
	(c)	(i) increases	1
		(ii) blue <i>ignore pale / dark etc</i>	1
		(iii) reversible (reaction) <i>allow goes both ways or two / either way</i>	1
		(iv) <u>anhydrous</u> copper sulfate	1

[8]

M5. (a) (i) the temperature at start  
*ignore reference to bubbles / heat* 1

the temperature at end  
*(measure) the temperature rise / change = 2 marks*  
*(measure) the temperature 1 mark* 1

(ii) temperature would increase  
*allow it gets hot(ter) / warm(er) or heat given off*  
*allow energy released / transferred* 1

(b) any **one** from:

- volume of acid  
*allow amount*  
*allow liquid*
- temperature of acid
- size of magnesium ribbon  
*allow volume / mass / amount*
- surface area of magnesium  
*ignore size of test tube and reference to water*

1

(c) (i) (Test tube) B 1

(ii) produces bubbles faster  
*accept more bubbles*

**or**  
faster rate of reaction  
*allow most reactive*

1

(d) The particles move faster

1

The particles collide more often

1

[8]

- M6. (a) (i) increase 1
- (ii) energy is given out to the surroundings 1
- (b) (i) NO 1  
*allow 2NO*  
*ignore nitrogen oxide*  
*do **not** allow equations*
- (ii) harmful / poisonous (owtte) 1  
*allow dangerous*  
*ignore reference to pollution / global warming*  
*do **not** accept references to ozone layer*
- (c) a catalyst can speed up a chemical reaction 1
- different reactions need different catalysts 1
- (d) (i) smaller 1  
*accept less / tiny / very small*  
*allow  $10^9$*   
*do **not** allow small unless qualified*
- (ii) reduce cost (owtte) **or**



*ignore references to energy*

save resources / raw materials (owtte)

1

[8]

M7. (a) (i) 4 1

(ii) (Make) 3 1

biggest temperature rise 1

(b) (i) 1008 (kJ)  
*correct answer with or without working gains 2 marks*  
*if incorrect answer given allow evidence of  $240 \times 4.2$  for 1 mark* 2

(ii) crisps have a high energy content  
*allow crisps have lots of calories / kilojoules / fat / one ninth of daily energy intake* 1

so if you take in more energy than you need the excess is stored as fat  
*accept consequences: obesity; heart disease; high blood pressure; diabetes; arthritis*

**or**

crisps contain salt (1)

too much salt can cause high blood pressure **or** heart problems or kidney problems (1)

1

[7]

- M8.** (a) goes up 1
- (b) (i) B 1
- (ii) A 1
- (iii) a catalyst 1
- activation energy 1
- (c) (i) eg (ensures) complete reaction  
*allow spread heat / energy*  
**or** even heating  
*allow mixes properly or mix them together or to get correct temperature*  
*ignore dissolves* 1
- (ii) lid (on beaker)  
*accept cover beaker*  
**or**  
insulate (beaker) / use a plastic cup 1

[7]

**Q1.**The figure below shows magnesium burning in air.



© Charles D Winters/Science Photo Library

(a) Look at the figure above.

How can you tell that a chemical reaction is taking place?

.....  
.....

(1)

(b) Name the product from the reaction of magnesium in the figure.

.....

(1)

(c) The magnesium needed heating before it would react.

What conclusion can you draw from this?

Tick **one** box.

The reaction is reversible

The reaction has a high activation energy

The reaction is exothermic

Magnesium has a high melting point

(1)

- (d) A sample of the product from the reaction in the figure above was added to water and shaken.

Universal indicator was added.

The universal indicator turned blue.

What is the pH value of the solution?

Tick **one** box.

1

4

7

9

(1)

- (e) Why are nanoparticles effective in very small quantities?

Tick **one** box.

They are elements

They are highly reactive

They have a low melting point

They have a high surface area to volume ratio

(1)

(f) Give **one** advantage of using nanoparticles in sun creams.

.....  
.....

(1)

(g) Give **one** disadvantage of using nanoparticles in sun creams.

.....  
.....

(1)

(h) A coarse particle has a diameter of  $1 \times 10^{-6}$  m.  
A nanoparticle has a diameter of  $1.6 \times 10^{-9}$  m.


Calculate how many times bigger the diameter of the coarse particle is than the diameter of the nanoparticle.

.....  
.....  
.....  
.....

(2)

(Total 9 marks)

**Q2.** A student investigates the energy released when zinc powder reacts with copper sulfate solution. The student uses the apparatus shown in **Figure 1**.



The student:

- measures 100 cm<sup>3</sup> copper sulfate solution into a beaker
- measures the temperature of the copper sulfate solution
- puts 1 g zinc powder into the beaker
- stirs the mixture with a thermometer
- measures the highest temperature.

The student's results were:

Starting temperature = 21 °C

Highest temperature = 32 °C

(a) (i) Calculate the change in temperature.

.....

Change in temperature = ..... °C

(1)

(ii) Calculate the energy released in the reaction.

Use the equation

$$\begin{array}{ccccccc} \text{energy released} & = & \text{volume of} & & & & \\ \text{in J} & & \text{solution} & \times & 4.2 & \times & \text{temperature change} \\ & & \text{in cm}^3 & & & & \text{in } ^\circ\text{C} \end{array}$$

.....

.....

Energy released = ..... J

(2)


(b) The reaction of zinc with copper sulfate is exothermic.

How can you tell from the student's results that the reaction is exothermic?

.....  
.....

(1)

(c) The energy diagram for the reaction is shown in **Figure 2**.



(i) How can you tell from the energy diagram that the reaction is exothermic?

.....  
.....

(1)

(ii) Which arrow shows the activation energy in **Figure 2**?

Tick (✓) **one** box.

**A**

**B**



**c**



**(1)**  
**(Total 6 marks)**


**Q3.** The following steps show how to use a type of glue.

**Step 1** Measure out equal amounts of the liquids from tubes **A** and **B**.




**Step 2** Mix the liquids to make the glue.

Put a thin layer of the glue onto each of the surfaces to be joined.



**Step 3** Put the pieces together and hold them with tape.



**Step 4** Leave the glue to set.

(a) When liquids **A** and **B** are mixed a chemical reaction takes place.

This reaction is *exothermic*.

What does *exothermic* mean?

.....  
 .....  
 .....  
 .....

(2)

(b) The time taken for the glue to set at different temperatures is given in the table below.

Temperature in°C	Time taken for the glue to set
20	3 days
60	6 hours
90	1 hour

(i) Use the correct answer from the box to complete each sentence.

decreases	increases	stays the same
-----------	-----------	----------------

When the temperature is increased the time taken for the glue to set

.....

When the temperature is increased the rate of the setting reaction

.....

(2)

(ii) Tick (✓) **two** reasons why an increase in temperature affects the rate of reaction.

Reason	Tick (✓)
It gives the particles more energy	
It increases the concentration of the particles	

It increases the surface area of the particles	
It makes the particles move faster	

(2)  
(Total 6 marks)

**Q4.** Hand warmers use chemical reactions.



(a) The table shows temperature changes for chemical reactions **A**, **B** and **C**.

Reaction	Starting temperature in °C	Final temperature in °C	Change in temperature in °C
<b>A</b>	18	25	+ 7
<b>B</b>	17	.....	+ 5
<b>C</b>	18	27	+ 9

What is the final temperature for reaction **B**? Write your answer in the table.

(1)

(b) (i) What name is given to reactions that heat the surroundings? .....

(1)

(ii) Which reaction, **A**, **B** or **C**, would be best to use in a hand warmer?


Reaction

Give a reason why you chose this reaction.


.....  
.....

(2)

(c) A student added water to some anhydrous copper sulfate.



The equation for the reaction is shown.



The student measured the temperature before and after the reaction.

(i) The measurements showed that this reaction can be used for a hand warmer.

Draw a ring around the correct answer to complete the sentence.

When water is added to anhydrous copper sulfate the temperature

of the mixture

increases.

decreases.

stays the same.

(1)

(ii) Anhydrous copper sulfate is white.

What colour is seen after water is added to the anhydrous copper sulfate?

.....

(1)

(iii) What does the symbol  $\rightleftharpoons$  mean?

.....

(1)

(iv) The student heated a tube containing hydrated copper sulfate.

Name the solid substance produced.


.....

(1)

(Total 8 marks)

**Q5.** A student investigated the reaction of magnesium with hydrochloric acid.

(a) A piece of magnesium was dropped into the hydrochloric acid.



Bubbles of gas were produced and the magnesium disappeared.

The reaction is exothermic.

(i) What measurements would the student make to show that the reaction is exothermic?

.....  
.....  
.....  
.....

(2)

(ii) How would these measurements show that the reaction is exothermic?

.....

(1)

The student investigated how changing the concentration of the hydrochloric acid affects this reaction.

Each test tube contained a different concentration of hydrochloric acid.



The diagrams show the results of this experiment.



**Test tube A**



**Test tube B**



**Test tube C**



**Test tube D**

(b) Suggest **one** control variable in this investigation.

.....  
.....

(1)

(c) (i) Which test tube, **A**, **B**, **C** or **D**, contained the greatest concentration of hydrochloric acid?

Test tube

(1)

(ii) Why did you choose this test tube?

.....  
.....

(1)

(d) The student predicted that if the temperature of the acid was increased the reaction would take place faster.

Tick (✓) **two** statements in the table which explain why.

Statement	Tick (✓)
The particles move faster	
The particles collide with less energy	
The particles collide more often	
The particles are bigger	


(2)  
(Total 8 marks)

**Q6.** Read the information about car engines.

Burning petrol in air is an exothermic reaction. This reaction is used in car engines.

When petrol burns it produces harmful substances such as nitrogen oxides and carbon monoxide.

A catalytic converter stops these harmful substances being released into the air.



(a) Draw a ring around the correct answer to complete each sentence.

(i) The exothermic reaction makes the temperature of the engine

decrease.  
increase.  
stay the same.


(1)

(ii) This is because during exothermic reactions


energy is taken in from the surroundings.  
energy is given out to the surroundings.  
there is no energy change.

(1)

- (b) The diagram shows a catalytic converter which removes harmful substances. The catalytic converter has two parts, **A** and **B**, which contain different catalysts.



- (i) The equation for the reaction that takes place in part **A** is:




Which **one** of the substances shown in the equation is a compound?

Give the formula of this compound.

.....

(1)

- (ii) The equation for the reaction that takes place in part **B** is:



Why is it important to stop carbon monoxide (CO) from being released into the air?

.....

.....

(1)

- (c) The table lists some statements about catalysts. Only **two** statements are correct.

Tick (✓) the **two** correct statements.

Statement	Tick (✓)

A catalyst can speed up a chemical reaction.	
A catalyst is used up in a chemical reaction.	
Different reactions need different catalysts.	
A catalyst does <b>not</b> change the rate of a chemical reaction.	

(2)

- (d) Modern catalytic converters contain nanosized particles of catalyst. Less catalyst is needed when nanosized catalyst particles are used.

- (i) Complete the sentence.

The size of nanosized particles is ..... than normal sized particles.

(1)

- (ii) The catalysts contain platinum.


Suggest why a manufacturer of catalytic converters would want to use less catalyst.

.....  
 .....

(1)

(Total 8 marks)

**Q7.** A student investigated the amount of energy released when four different makes of plain salted crisps were burned.



The following method was used for each make of plain salted crisp. The pieces of crisp were all the same size.

- The starting temperature of the water was measured.
- The piece of crisp was burned underneath the test tube.
- The final temperature of the water was measured.

(a) The results of the investigation are shown in the table.

	Make 1	Make 2	Make 3	Make 4
Final temperature of the water in °C	26	25	29	25
Starting temperature of the water in °C	19	20	20	21
Temperature rise of the water in °C	7	5	9	

(i) Calculate the temperature rise for **make 4**.

.....

Temperature rise = ..... °C

(1)

(ii) Which make of crisp, **1, 2, 3** or **4**, releases the most energy?

Make .....

Give a reason for your answer.

.....

.....

(2)

(b) The energy needed by a student is about 9000 kJ each day.

(i) One large bag of crisps states that the energy released by the crisps is 240 kcal.

Calculate the energy of this bag of crisps in kJ.

1 kcal = 4.2 kJ

.....

.....

Answer = ..... kJ

(2)

(ii) Eating too many crisps is thought to be bad for your health.

Use the information above and your knowledge to explain why.

.....

.....

.....


.....

(2)

(Total 7 marks)

**Q8.** Hydrogen peroxide decomposes slowly to give water and oxygen.

The reaction is *exothermic*.



(a) In an *exothermic* reaction, energy is given out.

Draw a ring around the correct answer to complete the sentence.

In an *exothermic* reaction, the temperature


goes down.

goes up.

stays the same.

(1)


(b) The energy level diagram for this reaction is shown below.




The energy changes, **A**, **B** and **C**, are shown on the diagram.

Use the diagram to help you answer these questions.



Which energy change, <b>A</b> , <b>B</b> or <b>C</b> , is the activation energy?		
--	---	--

(1)

Which energy change, <b>A</b> , <b>B</b> or <b>C</b> , shows that this reaction is exothermic?		
--	---	--

(1)

- (iii) Hydrogen peroxide decomposes quickly when a small amount of manganese(IV) oxide is added.

Draw a ring around the correct answer to complete each sentence.

Hydrogen peroxide decomposes quickly because

manganese(IV) oxide is

a catalyst.

an element.

a solid.

activation energy.


The manganese(IV) oxide has lowered the

boiling point.  
temperature.

(2)

- (c) A student did an experiment to find the amount of energy produced when hydrogen peroxide solution is decomposed using manganese(IV) oxide.

The apparatus the student used is shown in the diagram.



The student first measured the temperature of the hydrogen peroxide. Then the student added the manganese(IV) oxide, stirred the mixture and recorded the highest temperature.

- (i) Suggest why the student stirred the mixture before recording the highest temperature.

.....  
.....

(1)

- (ii) The biggest error in this experiment is heat loss.

Suggest how the student could change the apparatus so that less heat is lost.

.....  
.....

(1)

(Total 7 marks)

**M1.(a)** any **one** from:

- solution becomes colourless or colour fades
- zinc becomes bronze / copper coloured  
*allow copper (forms) or a solid (forms)*
- zinc gets smaller  
*allow zinc dissolves*
- bubbles or fizzing.  
*ignore precipitate*

1

(b) improvement:

use a plastic / polystyrene cup or add a lid

*accept use lagging / insulation*

1

reason - must be linked

reduce / stop heat loss

**OR**

improvement:

use a digital thermometer

*allow use a data logger*

reason - must be linked

more accurate or easy to read or stores data

*allow more precise or more sensitive*

*ignore more reliable*

*ignore improvements to method, eg take more readings*

1

(c) Marks awarded for this answer will be determined by the Quality of Written Communication (QWC) as well as the standard of the scientific response. Examiners should also refer to the information in the Marking Guidance and apply a 'best-fit' approach to the marking.

**0 marks**

No relevant content

**Level 1 (1–2 marks)**

There is a statement about the results.

**Level 2 (3–4 marks)**

There are statements about the results. These statements may be linked or may include data.

**Level 3 (5–6 marks)**

There are statements about the results with at least one link and an attempt at an explanation.

Examples of chemistry points made in the response:

**Description:****Statements**

Concentration of copper sulfate increases

Temperature change increases

There is an anomalous result

The temperature change levels off

Reaction is exothermic

**Linked Statements**

Temperature change increases as concentration of copper sulfate increases

The temperature change increases, and then remains constant

After experiment 7 the temperature change remains constant

**Statements including data**

The trend changes at experiment 7

Experiment 3 is anomalous

**Attempted Explanation**

Temperature change increases because rate increases

Temperature change levels off because the reaction is complete

**Explanation**

As more copper sulfate reacts, more heat energy is given off

Once copper sulfate is in excess, no further heat energy produced

6

[9]

**M2.(a)** any **three** from:

- concentration of (salt) solution
- volume of (salt) solution  
*ignore amount of solution*
- **initial** temperature (of the solution)  
*ignore room temperature*
- surface area / form of metal
- moles of metal  
*allow mass / amount*  
*ignore time*  
*ignore size of tube*

**3**

(b) 20

**1**

32

**1**

12

*allow ecf*

**1**

- (c) (i) four bars of correct height  
*tolerance is + / - half square*  
*3 correct for 1 mark*

**2**

bars labelled

**1**

- (ii) *one variable* is non-continuous / categoric  
*accept qualitative or discrete*

*accept no values between the metals*

1

(iii) magnesium

1

because biggest temperature change

*accept gives out most energy*

*ignore rate of reaction*

*dependent on first mark*

1

(iv) does not react / silver cannot displace copper

1

because silver not more reactive (than copper) **or** silver below copper in reactivity series

*do **not** accept silver is less reactive than copper sulfate*

1

(v) replace the copper sulfate

*could be implied*

1

with any compound of a named metal less reactive than copper

*allow students to score even if use an insoluble salt*

1

[16]

- M3.(a)** eg plastic (beaker) / insulation / lid / cover **or** any mention of enclosed  
*any sensible modification to reduce heat loss*  
*ignore prevent draughts*  
*ignore references to gas loss*  
*ignore bomb calorimeter* 1
- (b) all the substances react **or** all (the substances) react fully / completely **or** heat evolved quickly **or** distribute heat  
*'so they react' is insufficient for the mark*  
*accept increase chances of (successful) collisions / collision rate increase*  
*do **not** accept rate of reaction increase / make reaction faster* 1
- (c) experiment 2 **and**  
 different / higher / initial / starting temperature  
*accept experiment 2 **and** the room is hotter / at higher temperature*  
*do **not** accept temperature change / results higher* 1
- (d) temperature change does not fit pattern  
*accept anomalous / odd **or** it is the lowest **or** it is lower than the others **or** it is different to the others*  
*'results are different' is insufficient* 1
- (e) 7 / 7.0 1
- (f)  $(100 \times 4.2 \times 7) = 2940$   
*ecf from (e)* 1

- (g) diagram A **and**  
reaction exothermic / heat evolved /  $\Delta H$  is negative / temperature rises  
*accept energy is lost (to the surroundings)*  
*accept energy of products lower than reactants*  
*allow arrow goes downwards*

1

[7]



- M4.** (a) any **one** from:
- no method / electrolysis / equipment / technology  
*allow 'didn't know how to' or 'no knowledge'*
  - aluminium is a very reactive metal
  - high melting point  
*allow 'couldn't heat it enough'*
  - potassium had not been discovered
- 1**
- (b) because others / scientists / they could not repeat the experiment  
*ignore he could not repeat the experiment*
- or**  
others / they could not obtain the same results
- 1**
- (c) reaction is endothermic **or**  
reaction takes in heat / energy  
*accept activation energy*  
*ignore rate / high temperature*  
*ignore bonds broken*
- 1**
- (d) (aluminium chloride + potassium) → aluminium + potassium chloride  
*in either order*  
*accept correct formulae*  
*ignore metal*  
*ignore balancing*
- 1**
- (e) when tested it had the properties of a metal  
*accept a test for a metal property eg conductivity / reaction with acid*
- 1**

properties were different (from other known metals)  
*accept properties compared with other metals*

1

[6]

M5. (a) gives out energy **or** heat 1

(b) (i) *accept qualified answers in terms of volume of gas related to time*  
fast initially 1

slows down 1

reaction stops  
*accept reaction is now very slow* 1

(b) (ii) 21 1

(iii) 84  
*correct answer with or without working = 2 marks*  
*allow ecf from (b)(ii) correctly calculated for 2 marks*  
*allow evidence of 21/25 **or** (b)(ii)/25 for 1 mark* 2

(c) because they / particles have more energy / move faster  
*ignore particles move more / vibrate* 1

(and so) particles collide more often / more frequently **or** particles more likely to collide  
*ignore collide faster*  
*ignore more collisions* 1

(and) more of the collisions are successful **or** particles collide with more energy / harder **or** more of the particles have the activation energy

*accept more successful collisions*

1

[10]

**M6.** (a) gives out heat / energy  
*allow release / loses*  
*allow the products have less energy*

**or**

energy / heat transferred to the surroundings  
*ignore temperature rises*  
*allow more energy given out in forming bonds than taken in to break bonds*

1

(b) (i) speed up the reaction (owtte)  
*accept changes the rate*  
*accept lowers activation energy*  
*accept increases successful collisions*  
*accept allows reaction to take place at a lower temperature*

1

(ii) nitrogen (N<sub>2</sub>) / oxygen (O<sub>2</sub>) / products are safe **or** not harmful / pollutant / toxic / dangerous / damaging  
*ignore releases nitrogen / oxygen unless qualified*

**or**

(harmful) nitrogen monoxide / NO is not released into the air.  
*accept prevents / less acid rain*  
*ignore greenhouse gas / ozone layer*

1

(iii) 2 and 2  
*accept correct multiples or fractions*

1

(iv) idea of catalyst not being used up  
*allow not changed by reaction*  
*ignore catalyst does not take part*

*ignore catalyst not used in the reaction*

1

(v) idea of different reactions (require different catalysts)

*accept catalysts work for specific reactions*

*allow different gases*

1

(c) • smaller / very small / or any indication of very small / 1–100 nanometres /  
a few (hundred) atoms

*ignore just small*

*ignore size of the converter*

1

• big(ger) surface area

1

• less (catalyst) needed / small amount of catalyst needed


1

[9]

**Q1.** A student investigated the temperature change when zinc reacts with copper sulfate solution.

The student used a different concentration of copper sulfate solution for each experiment.

The student used the apparatus shown below.



The student:

- measured 50 cm<sup>3</sup> copper sulfate solution into a glass beaker
- measured the temperature of the copper sulfate solution
- added 2.3 g zinc
- measured the highest temperature
- repeated the experiment using copper sulfate solution with different concentrations.

The equation for the reaction is:



(a) The thermometer reading changes during the reaction.

Give **one** other change the student could **see** during the reaction.

.....  
.....

(1)

(b) Suggest **one** improvement the student could make to the apparatus.

Give a reason why this improves the investigation.

Improvement .....

.....  
Reason .....

.....

(2)

- (c) **In this question you will be assessed on using good English, organising information clearly and using specialist terms where appropriate.**

The student's results are shown in the table.

**Table**

<b>Experiment number</b>	<b>Concentration of copper sulfate in moles per dm<sup>3</sup></b>	<b>Increase in temperature in °C</b>
1	0.1	5
2	0.2	10
3	0.3	12
4	0.4	20
5	0.5	25
6	0.6	30
7	0.7	35
8	0.8	35
9	0.9	35
10	1.0	35

Describe **and** explain the trends shown in the student's results.

.....  
.....  
.....  
.....



.....

.....

.....

.....

.....

.....

(6)  
(Total 9 marks)


**Q2.** A student investigated displacement reactions of metals.

The student added different metals to copper sulfate solution and measured the temperature change.

The more reactive the metal is compared with copper, the bigger the temperature change.

The apparatus the student used is shown in **Figure 1**.

**Figure 1**



(a) State **three** variables that the student must control to make his investigation a fair test.

1 .....

2 .....

3 .....


**(3)**

(b) **Figure 2** shows the thermometer in one experiment before and after the student added a metal to the copper sulfate solution.

**Figure 2**

**Before adding metal**

**After adding metal**



Use **Figure 2** to complete **Table 1**.

**Table 1**

Temperature before adding metal in °C	.....
Temperature after adding metal in °C	.....
Change in temperature in °C	.....

(3)

- (c) The student repeated the experiment three times with each metal.

**Table 2** shows the mean temperature change for each metal.


**Table 2**

Metal	Mean temperature change in °C
Cobalt	4.5
Gold	0.0
Magnesium	10.0
Nickel	3.0
Silver	0.0

Tin	1.5
-----	-----

(i) On **Figure 3**, draw a bar chart to show the results.

**Figure 3**



(3)

(ii) Why is a line graph **not** a suitable way of showing the results?

.....  
 .....

(1)

(iii) Use the results to work out which metal is the most reactive.

Give a reason for your answer.

Most reactive metal .....

Reason .....

.....

(2)

(iv) Explain why there was no temperature change when silver metal was added to the copper sulfate solution.

.....  
.....  
.....  
.....

(2)

(v) It is **not** possible to put all six metals in order of reactivity using these results.

Suggest how you could change the experiment to be able to put all six metals into order of reactivity.

.....  
.....  
.....  
.....  
.....


(2)

(Total 16 marks)


**Q3.** Read the information about energy changes and then answer the questions.

A student did an experiment to find the energy change when hydrochloric acid reacts with sodium hydroxide.

The equation which represents the reaction is:



The student used the apparatus shown in the diagram.



The student placed 50 cm<sup>3</sup> of hydrochloric acid in a glass beaker and measured the initial temperature.

The student then quickly added 50 cm<sup>3</sup> of sodium hydroxide solution and stirred the mixture with the thermometer. The highest temperature was recorded.

The student repeated the experiment, and calculated the temperature change each time.

	Experimen t 1	Experimen t 2	Experimen t 3	Experimen t 4
Initial temperature in °C	19.0	22.0	19.2	19.0
Highest temperature in °C	26.2	29.0	26.0	23.5
Temperature change in °C	7.2	7.0	6.8	4.5

(a) The biggest error in this experiment is heat loss.

Suggest how the apparatus could be modified to reduce heat loss.

.....  
.....

(1)

(b) Suggest why it is important to mix the chemicals thoroughly.

.....

(1)

(c) Which **one** of these experiments was probably done on a different day to the others?

Give a reason for your answer.

.....

(1)

(d) Suggest why experiment **4** should **not** be used to calculate the average temperature change.

.....

.....

(1)

(e) Calculate the average temperature change from the first three experiments.

.....

Answer = ..... °C

(1)

(f) Use the following equation to calculate the energy change for this reaction.

$$\text{Energy change in joules} = 100 \times 4.2 \times \text{average temperature change}$$


.....

Answer = ..... J

(1)

(g) Which **one** of these energy level diagrams represents the energy change for this reaction?

Give a reason for your answer.



.....  
.....

(1)  
(Total 7 marks)



**Q4.** Read the information.

Alumina is a white solid. In 1800, scientists thought that alumina contained an undiscovered metal. We now call this metal aluminium. At that time, scientists could not extract the aluminium from alumina.

In 1825, Christian Oersted, a Danish scientist, did experiments with alumina.

**Step 1** He reacted a mixture of hot alumina and carbon with chlorine to form aluminium chloride. The reaction is very endothermic.

**Step 2** The aluminium chloride was reacted with potassium. He was left with potassium chloride and tiny particles of aluminium metal.

Other scientists were **not** able to obtain the same results using his experiment and his work was not accepted at that time.

In 1827, Friedrich Wöhler, a German chemist, made some changes to Oersted's experiment. He obtained a lump of aluminium. He tested the aluminium and recorded its properties.

(a) Suggest why scientists in 1800 could not extract aluminium from alumina.

.....  
.....

(1)

(b) Oersted's experiment in 1825 was **not** thought to be reliable.

Explain why

.....  
.....

(1)

(c) Why must the reaction in **Step 1** be heated to make it work?

.....  
.....

(1)

(d) Complete the word equation for the reaction in **Step 2**.

aluminium +potassiu→..... +.....  
chloride m

(1)


(e) Suggest how Wöhler was able to prove that he had made a new metal.

.....  
.....  
.....  
.....

(2)

(Total 6 marks)

**Q5.** The symbol equation for the decomposition of hydrogen peroxide is:




(a) This reaction is *exothermic*.

What is an *exothermic* reaction?


.....  
.....

(1)

(b) A student measured the volume of oxygen produced by 50 cm<sup>3</sup> of hydrogen peroxide.



The graph shows the results.



- (i) Use the graph to describe the changes in the rate of the reaction from 0 to 35 seconds.

.....

.....

.....

.....

.....

.....

(3)

- (ii) What was the total volume of oxygen gas collected?

..... cm<sup>3</sup>

(1)

- (iii) The student had calculated that the hydrogen peroxide used should produce 25 cm<sup>3</sup> of oxygen.

Calculate the percentage yield of oxygen.

.....

.....

.....

Answer = ..... %

(2)

(c) An increase in the temperature of the hydrogen peroxide increases the rate of the reaction.

Use your knowledge of particles to explain why.

.....  
.....  
.....  
.....  
.....  
.....

(3)


(Total 10 marks)

**Q6.** Read the information about car engines.

Burning petrol in air is an *exothermic* reaction. This reaction is used in car engines.

When petrol burns it produces harmful substances such as nitrogen oxides and carbon monoxide.

A catalytic converter stops these harmful substances being released into the air.




(a) The reaction is *exothermic*. What is the meaning of *exothermic*?

.....  
.....

(1)

(b) The catalytic converter has two parts shown as **A** and **B** in the diagram.



Part **A** contains a catalyst made from platinum and rhodium.

Part **B** contains a catalyst made from platinum and palladium.


(i) Why are catalysts used in chemical reactions?

.....

.....

(1)

(ii) One reaction in part **A** is shown by this equation.




Suggest why this reaction helps the environment.

.....  
.....

(1)

(iii) The equation for one of the reactions in part **B** is shown below.

Balance this equation.



(1)

(iv) The catalytic converter works for many years without replacing the catalyst.

Explain why the catalyst does not need to be replaced.

.....  
.....

(1)

(v) Suggest why different catalysts are used in parts **A** and **B**.

.....  
.....

(1)

- (c) Modern catalytic converters contain nanosized particles of catalyst. Using nanosized particles reduces the cost of the catalytic converter.

Suggest and explain why the use of nanosized catalyst particles reduces the cost of the catalytic converter.

Your answer should include information about the size and surface area of the particles.

.....

.....

.....

.....

.....

.....

.....

.....

**(3)**  
**(Total 9 marks)**



M1.(a) line goes up before it goes down 1

energy given out correctly labelled 1

activation energy labelled correctly 1

(b) electrostatic force of attraction between shared pair of negatively charged electrons 1

and both positively charged nuclei 1

(c) bonds formed =  $348 + 4(412) + 2(276) = 2548 \text{ kJ / mol}$  1

bonds broken – bonds formed =  $612 + 4(412) + (\text{Br-Br}) - 2548 = 95 \text{ kJ / mol}$  1

*Alternative approach without using C-H bonds*

*For step 1 allow =  $348 + 2(276) = 900 \text{ kJ / mol}$*

*Then for step 2 allow  $612 + (\text{Br-Br}) - 900 = 95 \text{ kJ / mol}$*

193 (kJ / mol) 1

*accept (+)193 (kJ / mol) with no working shown for 3 marks*

$-193(\text{kJ} / \text{mol})$  scores 2 marks  
allow ecf from step 1 and step 2

(d) **Level 3 (5–6 marks):**

A detailed and coherent explanation is given, which demonstrates a broad understanding of the key scientific ideas. The response makes logical links between the points raised and uses sufficient examples to support these links. A conclusion is reached.

**Level 2 (3–4 marks):**

An explanation is given which demonstrates a reasonable understanding of the key scientific ideas. A conclusion may be reached but the logic used may not be clear or linked to bond energies.

**Level 1 (1–2 marks):**

Simple statements are made which demonstrate a basic understanding of some of the relevant ideas. The response may fail to make logical links between the points raised.

**0 marks:**

No relevant content.

**Indicative content**

Size and strength

- chlorine atoms have fewer electron energy levels / shells
- chlorine atoms form stronger bonds
- Cl–Cl bond stronger than Br–Br
- C–Cl bond stronger than C–Br

Energies required

- more energy required to break bonds with chlorine
- more energy given out when making bonds with chlorine
- overall energy change depends on sizes of energy changes

Conclusions

- if C–Cl bond changes more, then less exothermic
- if C–Cl bond changes more then more exothermic
- can't tell how overall energy change will differ as do not know which changes more.

6

[14]

M2.(a) (i) the products are at a lower energy level than the reactants

*accept products have less energy / less energy at the end than the beginning*

1

(ii) because a catalyst provides an alternative / different pathway / mechanism / reaction route

*accept adsorption or 'increases concentration at the surface'*

*ignore absorption*

1

(that has) lower activation energy

*allow weakens bonds*

*allow idea of increased successful collisions.*

*DO NOT ALLOW answers stating catalysts provide energy for M1 and M2*


1

(b) one pair of electrons in each overlap (8 pairs in total)

*allow any combination of dots, crosses or other symbols*

1

the rest of the diagram correct with four non-bonding electrons on the oxygen giving a total of eight electrons in oxygen outer energy level.



*gains 2 marks*

1

(c) (i)  $\pm 3024$  (J)

*correct answer with or without working gains 3 marks*

*if the answer is incorrect, award up to 2 marks for the following steps:*

- $\Delta T = 14.4(^{\circ}\text{C})$
- $50 \times 4.2 \times 14.4$

*allow ecf for incorrect  $\Delta T$*

3

(ii) 0.015(2173913)

correct answer with or without working gains **3** marks

if answer is incorrect, allow 1 mark each for any of the following steps up to a max of 2.

- 0.70g
- $M_r$  of ethanol = 46
- $0.70 / 46$

allow ecf in final answer for arithmetical errors

**3**

(iii)  $\pm 198\,720$  (J / mole)

$c(i) \div c(ii)$

allow ecf from **(c)(i)** and **(c)(ii)**

0.015 gives 201600

0.0152 gives 198947

0.01522 gives 198686

**1**

(d) (as the molecules get bigger **or** the number of carbon atoms increases) the intermolecular forces

allow intermolecular bonds

**1**

(intermolecular forces) increase

allow more / stronger (intermolecular forces)

**1**

and therefore require more (heat) energy to overcome

breaking covalent bonds or unspecified bonds max **1** mark (M3)

**1**

**[15]**

M3.(a) 31

1

(b) (i) any **two** from:

- incorrect reading of thermometer / temperature
- incorrect measurement of volume of acid
- incorrect measurement of volume of alkali (burette).

2

(ii) glass is a (heat) conductor **or** polystyrene is a (heat) insulator

*answer needs to convey idea that heat lost using glass **or** not lost using polystyrene*

*accept answers based on greater thermal capacity of glass (such as "glass absorbs more heat than polystyrene")*

1

(c) (i) temperature increases

1

(ii) no reaction takes place **or** all acid used up **or** potassium hydroxide in excess

1

cool / colder potassium hydroxide absorbs energy **or** lowers temperature

*ignore idea of heat energy being lost to surroundings*

1

(iii) take more readings

*ignore just "repeat"*

1

around the turning point **or** between 20 cm<sup>3</sup> and 32 cm<sup>3</sup>

*accept smaller ranges as long as no lower than 20 cm<sup>3</sup> and no higher than 32 cm<sup>3</sup>*

1

(d) 1.61 **or** 1.6(12903)

*correct answer with or without working scores **3***

*if answer incorrect, allow a maximum of **two** from:*

*moles nitric acid =  $(2 \times 25 / 1000) = 0.05$  for **1** mark*

*moles KOH = (moles nitric acid) = 0.05 for **1** mark*

*concentration KOH = 0.05 / 0.031*

*answer must be correctly rounded (1.62 is incorrect)*

3

(e) same amount of energy given out

1

which is used to heat a smaller total volume **or** mixture has lower thermal capacity

**or**

number of moles reacting is the same

but the total volume / thermal capacity is less

*if no other marks awarded award 1 mark for idea of reacting faster*

1

[14]

M4.(a) circle round any one (or more) of the covalent bonds

*any correct indication of the bond – the line between letters*

1

(b) Methane contains atoms of two elements, combined chemically


1

(c) (i) activation energy labelled from level of reagents to highest point of curve

*ignore arrowheads*

1

enthalpy change labelled from reagents to products



*arrowhead **must** go from reagents to products only*

1

(ii) 2 O<sub>2</sub>

1

2 H<sub>2</sub>O

*if not fully correct, award 1 mark for all formulae correct.*

*ignore state symbols*

1

(iii) carbon monoxide is made

1

this combines with the blood / haemoglobin **or** prevents oxygen being carried in the blood / round body **or** kills you **or** is toxic **or** poisonous

*dependent on first marking point*

1

(iv) energy is taken in / required to break bonds

*accept bond breaking is endothermic*

1

energy is given out when bonds are made  
*accept bond making is exothermic*

1

the energy given out is greater than the energy taken in  
*this mark only awarded if both of previous marks awarded*

1

(d) (i) energy to break bonds = 1895  
*calculation with no explanation max = 2*

1

energy from making bonds = 1998

1

1895 - 1998 (= -103)

**or**

energy to break bonds = 656

energy from making bonds = 759

656 - 759 (= -103)

*allow:*

*bonds broken - bonds made =*

*413 + 243 - 327 - 432 = -103 for 3 marks.*

1

(ii) The C — Br bond is weaker than the C — Cl bond


1

[15]



**Q1.** This question is about the reaction of ethene and bromine.

The equation for the reaction is:




(a) Complete the reaction profile in **Figure 1**.

Draw labelled arrows to show:

- The energy given out ( $\Delta H$ )
- The activation energy.

**Figure 1**



(3)

(b) When ethene reacts with bromine, energy is required to break covalent bonds in the molecules.

Explain how a covalent bond holds two atoms together.

.....

.....


.....

.....

(2)

- (c) **Figure 2** shows the displayed formulae for the reaction of ethene with bromine.

**Figure 2**



The bond enthalpies and the overall energy change are shown in the table below.

	C=C	C-H	C-C	C-Br	Overall energy change
Energy in kJ / mole	612	412	348	276	-95

Use the information in the table above and **Figure 2** to calculate the bond energy for the Br-Br bond.


.....  
 .....  
 .....

Bond energy ..... kJ / mole

(3)

- (d) **Figure 3** shows the reaction between ethene and chlorine and is similar to the reaction between ethene and bromine.

**Figure 3**



“The more energy levels (shells) of electrons an atom has, the weaker the covalent bonds that it forms.”

Use the above statement to predict and explain how the overall energy change for the reaction of ethene with chlorine will differ from the overall energy change for the reaction

of ethene with bromine.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


(6)  
(Total 14 marks)

**Q2.** This question is about ethanol.

(a) Ethanol is produced by the reaction of ethene and steam:



(i) **Figure 1** shows the energy level diagram for the reaction.



How does the energy level diagram show that the reaction is exothermic?

.....  
.....

(1)

(ii) A catalyst is used for the reaction.


Explain how a catalyst increases the rate of the reaction.

.....  
.....  
.....  
.....

(2)

(b) **Figure 2** shows the displayed structure of ethanol.


**Figure 2**



Complete the dot and cross diagram in **Figure 3** to show the bonding in ethanol.

Show the outer shell electrons only.


**Figure 3**



(2)

- (c) A student burned some ethanol.

**Figure 4** shows the apparatus the student used.



- (i) The student recorded the temperature of the water before and after heating.

His results are shown in **Table 1**.

**Table 1**

Temperature before heating	20.7 °C
Temperature after heating	35.1 °C

Calculate the energy used to heat the water.

Use the equation  $Q = m \times c \times \Delta T$

The specific heat capacity of water = 4.2 J / g / °C

.....  
.....  
.....  
.....  
.....

Energy used = ..... J

(3)

- (ii) **Table 2** shows the mass of the spirit burner before the ethanol was burned and after the ethanol was burned.

**Table 2**

Mass of spirit burner before ethanol was burned	72.80 g
Mass of spirit burner after ethanol was burned	72.10 g

Calculate the number of moles of ethanol (C<sub>2</sub>H<sub>5</sub>OH) that were burned.

Relative atomic masses (A<sub>r</sub>): H = 1; C = 12; O = 16

.....  
.....  
.....  
.....

.....  
Number of moles burned = .....

(3)

(iii) Calculate the energy released in joules per mole.

You should assume that all the energy from the ethanol burning was used to heat the water.

.....  
Energy = ..... J / mole

(1)

(d) The names, structures and boiling points of ethanol and two other alcohols are shown in **Table 3**.

**Table 3**

Name	Methanol	Ethanol	Propanol
Structure	$\begin{array}{c} \text{H} \\   \\ \text{H}-\text{C}-\text{O}-\text{H} \\   \\ \text{H} \end{array}$	$\begin{array}{c} \text{H} \quad \text{H} \\   \quad   \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{H} \\   \quad   \\ \text{H} \quad \text{H} \end{array}$	$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\   \quad   \quad   \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{O}-\text{H} \\   \quad   \quad   \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$
Boiling point in °C	65	78	97

Use your knowledge of structure and bonding to suggest why the boiling points increase as the number of carbon atoms increases.

.....  
.....  
.....  
.....  
.....  
.....

.....

(3)  
(Total 15 marks)



**Q3.** Dilute nitric acid reacts with potassium hydroxide solution.

The equation for the reaction is:



A student investigated the temperature change in this reaction.


This is the method the student used.

- Step 1 Put 25 cm<sup>3</sup> of dilute nitric acid in a polystyrene cup.
- Step 2 Use a thermometer to measure the temperature of the dilute nitric acid.
- Step 3 Use a burette to add 4 cm<sup>3</sup> of potassium hydroxide solution to the dilute nitric acid and stir the mixture.
- Step 4 Use a thermometer to measure the highest temperature of the mixture.
- Step 5 Repeat steps 3 and 4 until 40 cm<sup>3</sup> of potassium hydroxide solution have been added.

The dilute nitric acid and the potassium hydroxide solution were both at room temperature.

- (a) **Figure 1** shows part of the thermometer after some potassium hydroxide solution had been added to the dilute nitric acid.

**Figure 1**



What is the temperature shown on the thermometer?

The temperature shown is ..... °C

(1)

- (b) Errors are possible in this experiment.

- (i) Suggest **two** causes of random error in the experiment.

.....

.....  
.....  
.....

(2)

(ii) Another student used a glass beaker instead of a polystyrene cup.

This caused a systematic error.


Why does using a glass beaker instead of a polystyrene cup cause a systematic error?

.....  
.....  
.....

(1)

(c) The results of the student using the polystyrene cup are shown in **Figure 2**.

**Figure 2**



(i) How do the results in **Figure 2** show that the reaction between dilute nitric acid and potassium hydroxide solution is exothermic?

.....  
.....

(1)

- (ii) Explain why the temperature readings decrease between 28 cm<sup>3</sup> and 40 cm<sup>3</sup> of potassium hydroxide solution added.

.....  
.....  
.....

(2)

- (iii) It is difficult to use the data in **Figure 2** to find the exact volume of potassium hydroxide solution that would give the maximum temperature.

Suggest further experimental work that the student should do to make it easier to find the exact volume of potassium hydroxide solution that would give the maximum temperature

.....  
.....  
.....  
.....

(2)

- (d) The student did further experimental work and found that 31.0 cm<sup>3</sup> of potassium hydroxide solution neutralised 25.0 cm<sup>3</sup> of dilute nitric acid.

The concentration of the dilute nitric acid was 2.0 moles per dm<sup>3</sup>.



Calculate the concentration of the potassium hydroxide solution in moles per dm<sup>3</sup>.

.....  
.....  
.....  
.....

.....  
.....

Concentration = ..... moles per dm<sup>3</sup>

(3)

- (e) The student repeated the original experiment using 25 cm<sup>3</sup> of dilute nitric acid in a polystyrene cup and potassium hydroxide solution that was twice the original concentration.

She found that:

- a smaller volume of potassium hydroxide solution was required to reach the maximum temperature
- the maximum temperature recorded was higher.

Explain why the maximum temperature recorded was higher.


.....  
.....  
.....  
.....

(2)

(Total 14 marks)

**Q4.** Methane (CH<sub>4</sub>) is used as a fuel.

(a) The displayed structure of methane is:



Draw a ring around a part of the displayed structure that represents a covalent bond.

(1)

(b) Why is methane a compound?

Tick (✓) **one** box.

Methane contains atoms of two elements, combined chemically.

Methane is not in the periodic table.

Methane is a mixture of two different elements.


(1)

(c) Methane burns in oxygen.

(i) The diagram below shows the energy level diagram for the complete combustion of methane.


Draw and label arrows on the diagram to show:

- the activation energy
- the enthalpy change,  $\Delta H$ .



(2)

(ii) Complete and balance the symbol equation for the complete combustion of methane.



(2)

(iii) Explain why the **incomplete** combustion of methane is dangerous.

.....

.....

.....

.....

(2)

(iv) Explain why, in terms of the energy involved in bond breaking and bond making, the combustion of methane is exothermic.

.....

.....

.....

.....

.....  
.....

(3)

(d) Methane reacts with chlorine in the presence of sunlight.

The equation for this reaction is:



Some bond dissociation energies are given in the table.


Bond	Bond dissociation energy in kJ per mole
C-H	413
C-Cl	327
Cl-Cl	243
H-Cl	432

(i) Show that the enthalpy change,  $\Delta H$ , for this reaction is  $-103$  kJ per mole.

.....  
.....  
.....  
.....  
.....  
.....

(3)

(ii) Methane also reacts with bromine in the presence of sunlight.



This reaction is less exothermic than the reaction between methane and chlorine.

The enthalpy change,  $\Delta H$ , is  $-45$  kJ per mole.

What is a possible reason for this?

Tick (✓) **one** box.

$\text{CH}_3\text{Br}$  has a lower boiling point than  $\text{CH}_3\text{Cl}$

The C-Br bond is weaker than the C-Cl bond.

The H-Cl bond is weaker than the H-Br bond.

Chlorine is more reactive than bromine.

(1)  
(Total 15 marks)



- M1.(a) 36 cm<sup>3</sup> 1
- (b) all points correct  
*± ½ small square* 2
- allow 1 mark if 6 or 7 of the points are correct*
- 2 best fit lines drawn  
*must not deviate towards anomalous point* 2
- allow 1 mark if 1 line correct*
- (c) The bung was not pushed in firmly enough. 1
- The measuring cylinder was not completely over the delivery tube. 1
- (d) as mass of lithium carbonate increases volume of gas produced increases 1
- linear / (directly) proportional 1
- (e) A gas / carbon dioxide is produced.  
*allow because the air in the tube expands* 1
- (f) any **one** from:

- Potassium carbonate does not decompose to produce carbon dioxide / a gas.
- Potassium carbonate does not decompose at the temperature of the Bunsen burner **or** the Bunsen burner is not hot enough to decompose potassium carbonate.
- When potassium carbonate decomposes a gas is not formed.

1

[11]

M2.(a) cotton wool 1

(b) all points correct 2  
 $\pm \frac{1}{2}$  small square

*allow 1 mark if 5 or 6 of the points are correct*

best fit line 1  
*must not deviate towards anomalous point*

(c) (mass) 1  
2.1 (g)  
*allow ecf from drawn best fit line*

(time) 1  
100 (s)

(d) a gas is produced 1

which escapes from the flask 1

(e)  $\frac{9.85}{150} = 0.0656$  1

0.07 (g / s)

*allow ecf answer correctly calculated to 2 decimal places*

1

(f) collect the gas in a gas syringe

1

measured the volume of gas

*allow carbon dioxide for gas*

1

*allow for 1 mark*

*collected gas*

**or**

*counted bubbles*

(g) The particles have more energy

1

The particles move faster

1

[14]

- M3.(a)** (i) oxygen, sulfur trioxide  
*both needed for mark* 1
- (ii) compound 1
- (b) increases  
*accept (goes) higher / (goes) up / (is) faster / (are) more frequent* 1
- (c) activation 1
- (d) catalyst **or** increase temperature 1
- [5]

**M4.(a)** heat / energy

1

given out / transfers to surroundings

*the mark for given out / transfers to cannot be awarded without  
heat / energy*

*allow given off*

1

(b) (i) decreases

1

increases

1

(ii) it gives the particles more energy

1

it makes the particles move faster

1

**[6]**

M5.(a) (i) In suntan creams 1

(ii) Much smaller 1

(b) (i) have a high surface area to volume ratio 1

(ii) because a catalyst provides an alternative / different pathway / mechanism / reaction route  
*accept adsorption or 'increases concentration at the surface'*  
*ignore absorption* 1

(that has) lower activation energy  
*allow weakens bonds*  
*allow idea of increased successful collisions*  
*max 1 mark for incorrect chemistry eg increased energy of particles* 1

[5]

<b>M6.</b>	(a)	(i)	10	1
		(ii)	OH <sup>-</sup>	1
	(b)	(i)	air	1
		(ii)	particles move faster	1
			particles collide more often	1
		(iii)	catalyst(s)	1
	(c)		liquid	1

[7]




**Q1.** Lithium carbonate reacts with dilute hydrochloric acid.

A group of students investigated the volume of gas produced.


This is the method used.

1. Place a known mass of lithium carbonate in a conical flask.
2. Measure  $10\text{ cm}^3$  of dilute hydrochloric acid using a measuring cylinder.
3. Pour the acid into the conical flask.
4. Place a bung in the flask and collect the gas as shown in **Figure 1**.



(a) **Figure 2** shows the measuring cylinder.

**Figure 2**



What volume of gas has been collected?

Volume = ..... cm<sup>3</sup>

(1)


(b) The table below shows the students' results.

Mass of lithium carbonate in g	Volume of gas in cm <sup>3</sup>
0.0	0
0.1	22
0.2	44
0.3	50
0.4	88
0.5	96
0.6	96
0.7	96

On **Figure 3**:

- Plot these results on the grid.
- Complete the graph by drawing **two** straight lines of best fit.

**Figure 3**



(4)

(c) What are **two** possible reasons for the anomalous result?

Tick **two** boxes.

Too much lithium carbonate was added.

The bung was not pushed in firmly enough.

There was too much water in the trough.

The measuring cylinder was not completely over the delivery

The conical flask was too small.

(2)

(d) Describe the pattern the graph shows up to 0.4 g of lithium carbonate added.

.....

.....


.....

.....


(2)

(e) Lithium carbonate decomposes when heated.

The equation shows the decomposition of lithium carbonate.



**Figure 4** shows the apparatus a student used to decompose lithium carbonate.



Why does the limewater bubble?

.....

.....

(1)

- (f) The student repeated the experiment with potassium carbonate.  
The limewater did not bubble.

Suggest why there were **no** bubbles in the limewater.

.....


.....

(1)

(Total 11 marks)

**Q2.** A student investigated the rate of reaction between marble chips and hydrochloric acid.

**Figure 1** shows the apparatus the student used.



(a) What is **A**?

Tick **one** box.

cotton wool

limestone

poly(ethene)

rubber bung

(1)

(b) **Table 1** shows the student's results for one investigation.

**Table 1**


Time in s	Mass lost in g
0	0.0
20	1.6
40	2.6
60	2.9

80	3.7
100	4.0
120	4.0

On **Figure 2**:

- Plot these results on the grid.
- Draw a line of best fit.

**Figure 2**



(3)

(c) Use **Figure 2** to complete **Table 2**.

**Table 2**

Mass lost after 0.5 minutes	..... g
Time taken to complete the reaction	..... s

(2)

(d) The equation for the reaction is:



Explain why there is a loss in mass in this investigation.

.....  
.....  
.....  
.....

(2)

(e) Another student investigated the rate of a different reaction.

**Table 3** shows the results from the different reaction.

**Table 3**

Mass lost when the reaction was complete	9.85 g
Time taken to complete the reaction	2 minutes 30 seconds

Calculate the mean rate of the reaction using **Table 3** and the equation:

$$\text{mean rate of reaction} = \frac{\text{mass lost in g}}{\text{time taken in s}}$$

Give your answer to two decimal places.

.....  
.....

Mean rate of reaction = ..... g / s

(2)

(f) The student measured the change in mass of the reactants.

Describe another method, other than measuring the change in mass of the reactions, that the student could have used to find the rate of the reaction between marble chips and



hydrochloric acid.

.....

.....

.....

.....

(2)

- (g) Another student planned to investigate the effect of temperature on the rate of reaction. The student predicted that the rate of reaction would increase as the temperature was increased.

Give **two** reasons why the student's prediction is correct.

Tick **two** boxes.

The particles are more concentrated.

The particles have a greater mass.


The particles have a larger surface area.

The particles have more energy.

The particles move faster.

(2)  
(Total 14 marks)

**Q3.(a)** The figure below represents the reaction of sulfur dioxide with oxygen.



(i) Complete the word equation for the reaction of sulfur dioxide with oxygen.

sulfur dioxide + ..... → .....

(1)

(ii) Draw a ring around the correct answer to complete the sentence.

Sulfur dioxide (SO<sub>2</sub>) is

- a compound.
- an element.
- a mixture.

(1)

(b) The reactants are gases.

When the pressure of the gases is increased, the reaction gets faster.

Complete the sentence.

When the pressure of the gases is increased,

the frequency of the collisions .....

(1)

(c) The particles need energy to react.

Complete the sentence.

The minimum amount of energy that particles need to react is called

the ..... energy.

(1)

(d) Give **one** way of increasing the rate of the reaction other than changing the pressure.

.....  
.....

(1)  
(Total 5 marks)


**Q4.** The following steps show how to use a type of glue.

**Step 1** Measure out equal amounts of the liquids from tubes **A** and **B**.




**Step 2** Mix the liquids to make the glue.

Put a thin layer of the glue onto each of the surfaces to be joined.



**Step 3** Put the pieces together and hold them with tape.



**Step 4** Leave the glue to set.

(a) When liquids **A** and **B** are mixed a chemical reaction takes place.

This reaction is *exothermic*.

What does *exothermic* mean?

.....  
.....  
.....  
.....

(2)

(b) The time taken for the glue to set at different temperatures is given in the table below.

Temperature in°C	Time taken for the glue to set
20	3 days
60	6 hours
90	1 hour

(i) Use the correct answer from the box to complete each sentence.

<b>decreases</b>	<b>increases</b>	<b>stays the same</b>
------------------	------------------	-----------------------

When the temperature is increased the time taken for the glue to set

.....

When the temperature is increased the rate of the setting reaction

.....

(2)

(ii) Tick (✓) **two** reasons why an increase in temperature affects the rate of reaction.

Reason	Tick (✓)
It gives the particles more energy	
It increases the concentration of the particles	

It increases the surface area of the particles	
It makes the particles move faster	

(2)  
(Total 6 marks)

**Q5.** Nanoparticles have many uses.

(a) (i) Tick (✓) **one** use of nanoparticles.

In the extraction of iron

In suntan creams

In the test for oxygen

(1)

(ii) How is the size of nanoparticles different from normal-sized particles?

Draw a ring around the correct answer.

**much smaller**

**same size**

**much larger**

(1)

(b) Very small amounts of cerium oxide nanoparticles can be added to diesel fuel.

The cerium oxide is a catalyst.

(i) Draw a ring around the correct answer to complete the sentence.

Only a very small amount of cerium oxide nanoparticles is needed because

the nanoparticles

are elements.

are very reactive.

have a high surface area to volume ratio.

(1)

(ii) Explain how a catalyst increases the rate of a reaction.

.....

.....

.....

.....

.....

(2)  
(Total 5 marks)



Q6. (a) Ammonia solution is used in cleaning products to remove grease from kitchen surfaces.



Ammonia solution is alkaline.

(i) Draw a ring around the number most likely to be the pH of ammonia solution.

**1**      **3**      **7**      **10**


(1)

(ii) Draw a ring around the ion in ammonia solution which makes it alkaline.

**Cl<sup>-</sup>**      **H<sup>+</sup>**      **Na<sup>+</sup>**      **OH<sup>-</sup>**

(1)

(b) Ammonia is made using the Haber process.



(i) Where does the nitrogen used in the Haber process come from?

Draw a ring around your answer.

**air**                      **natural gas**                      **water**

(1)

(ii) A high temperature of 450 °C is used in the reactor.

Tick (✓) **two** reasons in the table which explain why high temperatures make reactions faster.

Reasons	Tick (✓)
Particles move faster	
Particles are closer together	
Particles collide more often	
Particles have less energy	

(2)

(iii) The iron in the reactor speeds up the reaction but is not used up.

What is the name given to substances that speed up the chemical reaction but which are not used up during the reaction?

.....

(1)

(c) Complete the sentence.

The condenser separates the ammonia from the unreacted nitrogen and hydrogen by turning the ammonia into a .....

(1)

(Total 7 marks)

M1.(a) sulfur / sulphur / S / S(s)

1

(b) as the temperature increases, the rate of reaction increases

*allow two correct values for rate quoted (from graph) at different temperatures*

1

the rate of increase increases **or** there is an exponential relationship

*accept the rate of reaction increases slowly (from 20 °C to 50 °C) then increases more rapidly for 2 marks*

*answer MUST be based on rate / speed of reaction*

1

(c) (i) any **two** from:

- temperature (of the reactants)
- concentration of hydrochloric acid
- volume of hydrochloric acid
- volume of sodium thiosulfate
- the (size / darkness / thickness of the) cross
- total volume of solution.

*if no other marks gained, allow 1 mark for:  
rate of stirring*

**OR**

*amount of hydrochloric acid / sodium thiosulfate*

**OR**

*volume of solution*

2

(ii) (because as the concentration increases) the number of particles per unit volume increases **or** particles are closer together.

*idea of more particles in a given space is required for the first mark.*

*ignore references to area.*

1

(therefore) the frequency of (successful) collisions increases

*allow increased chance / probability of collisions*

*number of collisions increases is insufficient here.*

**must** mention per unit time or frequency.

*ignore speed of collisions.*

*if reference to space and time missing from M1 and M2 but they are otherwise correct, then award 1 mark.*

1

so the number of particles (per unit volume) doubles or (the frequency of collisions) doubles.

*students can score 2 marks for a qualitative explanation; the third mark is for a quantitative explanation.*

1

[8]

- M2.(a) (i) the higher the temperature, the greater the rate  
**or**  
 at 40 °C rate is faster than at 20 °C  
*accept the higher the temperature, the faster the reaction* 1
- (ii) 40 °C curve is steeper  
*accept the 40 °C line becomes horizontal sooner*  
*accept at higher temperatures the reaction finishes sooner*  
*accept reaction finishes sooner at 40 °C*  
*accept at higher temperatures the gas is produced faster*  
**or**  
 correct comparison of data from the graph 1
- (iii) 2 1
- (b) (i) Concentration of acid  
 Mass of marble chips 2
- (ii) increases rate  
*incorrect reference to energy = max 1* 1
- (because of) more frequent collisions (between particles)  
*accept particles are more likely to collide*  
*ignore more collisions*  
*ignore more successful collisions* 1
- (c) any **one** from:  
 • increases rate of reaction  
 • reduces energy required  
 • lower temperature can be used  
 • catalyst is not used up. 1

[8]

M3.(a) (i) precipitation

1

(ii) (aq) on left hand side

1

(s) on right hand side

1

(iii) potassium iodide

1

potassium nitrate

1

(iv) filtration

1

(b) (i) diffusion

1

(ii) iodide ions move / diffuse faster than lead ions **or** travel further in the same time

*Must be a comparison*

*Accept converse*

1

because the lead iodide forms much closer to the lead nitrate (or **X**) than the potassium iodide (or **Y**).

*allow because iodide ions are smaller than lead ions*

*allow references to potassium iodide and lead nitrate*

1

(iii) the particles / ions move / diffuse faster  
*ignore which particles / ions the student refers to*

1

because they have more energy **or** will collide / meet sooner  
*ignore reference to frequency of collisions*

1

[11]



M4.(a) time from when the heating is started until 1

the limewater turns cloudy / milky 1

(b) (i) the temperature was not high enough  
*accept the copper carbonate had not started to decompose / react*  
*accept it takes time to heat up the copper carbonate* 1

the bubbles of gas were air  
*accept no carbon dioxide produced* 1

(ii) the copper carbonate was decomposing / reacting  
*accept the temperature was high enough to cause decomposition*  
*/ a reaction* 1

so carbon dioxide was produced  
*allow correct word / symbol equation* 1

(iii) copper oxide was produced  
*allow correct word / symbol equation* 1

because the copper carbonate had completely decomposed / reacted  
*ignore all of the carbon dioxide had been given off* 1

[8]

M5.(a) because sulfur / S forms

1

which is insoluble / a solid / a precipitate

1

(b) (i) 32

*correct answer with or without working gains 2 marks*

*accept evidence of 31 + 33 / 2 for 1 mark*

*allow 35 for 1 mark*

2

(ii) reaction rate increases

*if incorrect reference to energy = max 2*

1

because of more particles (per unit volume)

*allow because particles are closer together*

1

and because there is an increase in frequency of collisions

*accept because particles are more likely to collide or higher chance of collision*

*ignore more (successful) collisions*

1

[7]

M6.(a) (i) a continuous straight line missing anomalous point  
*allow a line which does not start at zero / origin*

1

(ii) any **two** sensible errors eg  
*ignore systematic / zero error / weighing error **or** error unqualified*

- timing errors and / or example
- measurement errors and / or example
- apparatus errors and / or example
- human / experimental / reading / random error and / or example  
or 'did not do it right'  
*could be two from **same** category  
eg two timing errors – watch not started at the same time plus  
difficulty in deciding when the cross has disappeared.*
- temperature fluctuation
- anomalous point  
*accept outlier / wrong result*
- results not recorded correctly
- plotting error
- rate calculated incorrectly  
*ignore 'not repeated'*

2

(b) (i) straight line  
*allow as concentration increases the rate goes up **or** converse  
allow numerical example  
allow positive correlation  
allow same gradient  
ignore 'most points near / on line of best fit'*

1

(ii) because of an increase in frequency of collisions

*max 1 if incorrect reference to energy or if subatomic particle specified*

*accept because particles are more likely to collide or higher chance of collision*

*ignore more (successful) collisions*

1

because there are more particles (per unit volume)

*allow because particles are closer together*

1

[6]

**M7.** (a) gives out energy **or** heat 1

(b) (i) *accept qualified answers in terms of volume of gas related to time*  
fast initially 1

slows down 1

reaction stops  
*accept reaction is now very slow* 1

(b) (ii) 21 1

(iii) 84  
*correct answer with or without working = 2 marks*  
*allow ecf from (b)(ii) correctly calculated for 2 marks*  
*allow evidence of 21/25 **or** (b)(ii)/25 for 1 mark* 2

(c) because they / particles have more energy / move faster  
*ignore particles move more / vibrate* 1

(and so) particles collide more often / more frequently **or** particles more likely to collide  
*ignore collide faster*  
*ignore more collisions* 1

(and) more of the collisions are successful **or** particles collide with more energy / harder **or** more of the particles have the activation energy


*accept more successful collisions*

1

[10]

**Q1.** A student investigated the rate of reaction between sodium thiosulfate solution and dilute hydrochloric acid, as shown in **Figure 1**.

**Figure 1**



The reaction produced a precipitate, which made the mixture turn cloudy.

The student timed how long it took until she could no longer see the cross.

She calculated the rate of the reaction.

(a) The equation for the reaction is:



Name the product that made the mixture go cloudy.


.....

(1)

(b) The student investigated the effect of changing the temperature of the sodium thiosulfate solution on the rate of reaction.

She plotted her results on a graph, as shown in **Figure 2**.

Figure 2



Describe the trends shown in the student's results.

.....

.....

.....

.....

.....

(2)

(c) The student then investigated the effect of changing the concentration of sodium thiosulfate solution on the rate of the reaction.

(i) Suggest **two** variables the student would need to control to make sure that her results were valid.

.....



.....  
.....  
.....

(2)

(ii) From this investigation the student correctly concluded:

‘As the concentration of sodium thiosulfate solution doubles, the rate of reaction doubles.’

Explain the student’s conclusion in terms of particles.


.....  
.....  
.....  
.....  
.....  
.....  
.....

(3)

(Total 8 marks)

**Q2.** A student investigated the rate of reaction between calcium carbonate (marble chips) and hydrochloric acid.

The student used the apparatus shown in **Figure 1**.



The student:


- recorded the volume of gas collected every 5 seconds
- repeated the experiment using hydrochloric acid at different temperatures.

The equation for the reaction is:



(a) The student plotted results for the hydrochloric acid at 20 °C and 40 °C on a graph.

**Figure 2** shows the student's graph.



Use information from **Figure 2** to answer these questions.

- (i) State **one** conclusion the student could make about the effect of temperature on the rate of the reaction.

.....  
.....

(1)

- (ii) Give **one** reason why the student could make this conclusion.

.....  
.....

(1)

- (iii) For the hydrochloric acid at 60 °C the student had collected 30 cm<sup>3</sup> after 15 seconds.

Calculate the average rate of reaction from 0 to 15 seconds.

.....  
.....

Rate of reaction = ..... cm<sup>3</sup> per second

(1)

- (b) The student then investigated how the surface area of marble chips affected the rate of reaction.

- (i) Which **two** variables should the student keep constant?

Tick (✓) **two** boxes.

Amount of water in the trough

Concentration of acid

Mass of marble chips

Size of marble chips

Volume of measuring cylinder

(2)

(ii) Explain, in terms of particles and collisions, the effect that increasing the surface area of the marble chips has on the rate of reaction.

.....

.....

.....

.....

(2)

(c) Calcium carbonate is a catalyst for the industrial production of biodiesel.

Give **one** reason why using a catalyst reduces costs.

.....

.....

(1)

(Total 8 marks)

**Q3.**Lead nitrate solution reacts with potassium iodide solution.

The reaction produces a solid.

**Figure 1** shows the reaction occurring.

**Figure 1**



Lead Iodide By Der Kreole (own work) (CC-BY-3.0) via Wikimedia Commons

(a) (i) Give the name of this type of reaction.

Tick (✓) **one** box.

Combustion

Neutralisation

Precipitation


(1)

(ii) Write the missing state symbols in the chemical equation.



(2)

(iii) Complete the word equation for the reaction.



(2)

(iv) How is solid lead iodide separated from the solution?

Draw a ring around the correct answer.

**Distillation                  Electrolysis                  Filtration**


(1)

(b) A group of students investigated the movement of particles.

The students filled a container with water.


The students added a crystal of lead nitrate at position **X** and a crystal of potassium iodide at position **Y**, as shown in **Figure 2**.

**Figure 2 – view from above**



After 3 minutes solid lead iodide started to form at the position shown in **Figure 3**.

**Figure 3 – view from above**



(i) Tick (✓) the correct box to complete the sentence.

Lead ions and iodide ions move through the water by

diffusion.

evaporation.

neutralisation.

(1)

- (ii) What conclusion can you make about the speed of movement of lead ions compared with iodide ions?

Give a reason for your answer.

.....

.....

.....

.....

(2)

- (iii) The students repeated the experiment at a higher temperature.

The solid lead iodide formed after a shorter period of time.

Explain why, in terms of particles.

.....

.....

.....


.....

(2)

(Total 11 marks)

**Q4.**Carbon dioxide is produced when copper carbonate is heated.

A student investigated heating copper carbonate.  
 The student used the apparatus to measure how long it took for carbon dioxide to be produced.  
 The student also noted what happened during each minute for three minutes.



- (a) The student used changes to the limewater to measure how long it took for carbon dioxide to be produced.

Describe how.

.....

.....

.....

.....

(2)

- (b) The student wrote down her observations.

Time interval in minutes	Observations
Between 0 and 1	A slow release of gas bubbles. The limewater did not change. The solid in the test tube was green.
Between 1 and 2	A fast release of gas bubbles.



	The limewater changed at 1 minute 10 seconds.
Between 2 and 3	No release of gas bubbles. The solid in the test tube was black.

(i) Suggest the reason for the student's observations between 0 and 1 minute.

.....

.....

.....

.....

.....

(2)

(ii) Explain the student's observations between 1 and 2 minutes.

.....

.....

.....

.....

(2)

(iii) Explain the student's observations between 2 and 3 minutes.

.....

.....

.....

.....

(2)


**Q5.** A student investigated the rate of reaction between sodium thiosulfate and dilute hydrochloric acid.

The student placed a conical flask over a cross on a piece of paper.


The student mixed the solutions in the flask.

The solution slowly went cloudy.

The student timed how long it took until the cross could not be seen.



The equation for the reaction is:



(a) Explain why the solution goes cloudy.

.....

.....

.....

.....

(2)

(b) The student repeated the experiment with different concentrations of sodium thiosulfate.

Concentration of	Time taken until the cross could not be
------------------	---

sodium thiosulfate in moles per dm <sup>3</sup>	seen in seconds			
	Trial 1	Trial 2	Trial 3	Mean
0.040	71	67	69	69
0.060	42	45	45	44
0.080	31	41	33	

(i) Calculate the mean time for 0.080 moles per dm<sup>3</sup> of sodium thiosulfate.

.....  
 .....  
 .....

Mean = ..... seconds

(2)

(ii) Describe and explain, in terms of particles and collisions, the effect that increasing the concentration of sodium thiosulfate has on the rate of the reaction.


.....  
 .....  
 .....  
 .....  
 .....  
 .....

(3)

(Total 7 marks)

**Q6.** Sodium thiosulfate solution reacts with hydrochloric acid. As the reaction takes place the solution slowly turns cloudy.

The diagram shows a method of measuring the rate of this reaction.



A student used this method to study how changing the concentration of the sodium thiosulfate solution alters the rate of this reaction.

The student used different concentrations of sodium thiosulfate solution. All the other variables were kept the same.

The results of the experiments are shown on the graph below.


(a) (i) Draw a line of best fit on the graph.

(1)

(ii) Suggest **two** reasons why all of the points do not lie on the line of best fit.

.....  
.....  
.....  
.....  
.....

(2)



(b) (i) In a conclusion to the experiment the student stated that:

‘The rate of this reaction is directly proportional to the concentration of the sodium thiosulfate.’

How does the graph support this conclusion?

.....  
 .....

(1)


(ii) Explain, in terms of particles, why the rate of reaction increases when the concentration of sodium thiosulfate is increased.

.....  
 .....

.....

(2)  
(Total 6 marks)

**Q7.** The symbol equation for the decomposition of hydrogen peroxide is:



(a) This reaction is *exothermic*.


What is an *exothermic* reaction?

.....


.....

(1)

(b) A student measured the volume of oxygen produced by 50 cm<sup>3</sup> of hydrogen peroxide.



The graph shows the results.



- (i) Use the graph to describe the changes in the rate of the reaction from 0 to 35 seconds.

.....

.....

.....

.....

.....

.....

(3)

- (ii) What was the total volume of oxygen gas collected?

..... cm<sup>3</sup>

(1)

- (iii) The student had calculated that the hydrogen peroxide used should produce 25 cm<sup>3</sup> of oxygen.

Calculate the percentage yield of oxygen.

.....

.....



.....

Answer = ..... %

(2)

- (c) An increase in the temperature of the hydrogen peroxide increases the rate of the reaction.  
Use your knowledge of particles to explain why.

.....  
.....  
.....  
.....  
.....  
.....

(3)

(Total 10 marks)

**M1.(a)**  $\text{CaCO}_3 + 2\text{HCl} \rightarrow \text{CaCl}_2 + \text{H}_2\text{O} + \text{CO}_2$  **2**

*allow 1 mark for correct formulae*

(b) sensible scales, using at least half the grid for the points **1**

all points correct

*± ½ small square*

*allow 1 mark if 8 or 9 of the points are correct*

**2**

best fit line

**1**

(c) steeper line to left of original **1**

line finishes at same overall volume of gas collected

**1**

(d) acid particles used up **1**

*allow marble / reactant used up*

so concentration decreases

*allow surface area of marble decreases*

**1**

so less frequent collisions / fewer collisions per second  
*do not accept fewer collisions unqualified*

1

so rate decreases / reaction slows down

1

(e) mass lost of 2.2 (g)

1

time taken of  
270 s

*allow values in range 265 – 270*

1

$$\frac{2.2}{270} = 0.00814814$$

*allow ecf for values given for mass and time*

1

0.00815 (g / s)

**or**

$8.15 \times 10^{-3}$

*allow 1 mark for correct calculation of value to 3 sig figs  
accept 0.00815 or  $8.15 \times 10^{-3}$  with no working shown for 4 marks*

1

(f) correct tangent

1

eg 0.35 / 50

1

0.007

*allow values in range of 0.0065 – 0.0075*

**1**

$7 \times 10^{-3}$

**1**

*accept  $7 \times 10^{-3}$  with no working shown for 4 marks*

**[20]**

**M2.(a)** (s) (aq) (aq) (g)

*must be in this order*

**2 marks if all four correct**

**1 mark if 2 or 3 correct**

**2**

(b) (i) 55

*ignore units*

**1**

(ii) 54

*allow ecf from (b)(i)*

**1**

(iii) 0.92

*correct answer with or without working gains **2 marks***

*ecf from volume in (b)(i)*

*accept 2 d.p. up to calculator value*

*if answer incorrect, allow rate = (b)(i) / 60 for **1 mark***

**2**

(c) (i) circle round point at (48,22)

**1**

(ii) problem (1) and explanation (1)

*explanation **must** give lower volume of gas or slower reaction*

*ignore human error unless qualified*

**problem with bung**

e.g. bung not placed in firmly / quickly enough

so gas lost

**or**

**problem with reagent**

e.g. acid was diluted **or** acid not replaced

so reaction slower

**or**

**problem with temperature**

e.g. temperature was lower than recorded temperature

so reaction slower

**or**

**problem with measurement**

e.g. length of magnesium less than 8 cm **or** timed for less than a minute

so less gas produced

2

(d) repeat the experiment (several times)

1

because anomalous results could be excluded

1

and then the mean can be determined / calculated

*accept suggestion of alteration to method, which is explained as to why it would reduce the error, for 3 marks (e.g. place the magnesium in a container within the flask (1) so it can be tipped into the acid once the bung is in place (1). This will prevent anomalous results or gas loss (1))*

*ignore idea of more accurate gas syringe*

*ignore shorter time intervals*

1

(e) (i) use clean magnesium **or** use magnesium without oxide coating

1

compare results

1

(ii) **either**

measure the temperature of the acid before (adding magnesium)

1

and after adding magnesium

**or**

place the conical flask in a water bath (at 40 °C) (1)

compare results (1)

1

[16]

M3. (a) 118 1

(b) it loses / transfers electrons  
*it = Au / gold atom* 1

three electrons  
*sharing / covalency = max 1 mark* 1

(c) (i) O<sub>2</sub> 1

2 CO and 2 CO<sub>2</sub>  
or  
correct balancing of equation from O  
*accept correct multiples / fractions throughout* 1

(ii) *reference to incorrect bonding = 1 mark max*  
because carbon dioxide is simple molecular / small molecules 1

there are intermolecular forces (between the molecules)  
*allow intermolecular bonds* 1

so a small amount of energy needed (to separate molecules) or (*intermolecular forces*) are weak 1



(d) any **three** from:

- gold is the only catalyst for some reactions
- catalysts are not used up
- improves speed of reaction

reduces amount of energy **or** process needs low(er) temperature

*if no mark awarded, allow catalyst reduce costs (of the process)  
for 1 mark*

- only small quantities (of catalyst) needed

3

[11]

**M4.** (a) same number of (gaseous) molecules / moles / volume on both sides of the equation

*allow particles for molecules*

*do **not** accept atoms*

*ignore amount*

1

(b) (forward) reaction is exothermic

*accept reverse answer*

1

(c) any **three** from:

- particles gain energy
- particles move faster  
*allow particles collide faster / quicker*  
*ignore move more / vibrate more*
- particles collide more **or** more collisions
- more of the collisions are successful **or**  
more of the particles have the activation energy **or**  
particles collide with more force / energy

3

(d) any **two** from:

- more product (obtained in shorter time)  
*accept better yield (of product)*
- less fuel needed  
*accept less energy / heat / electricity needed*  
**or**  
lower fuel costs  
*ignore cheaper unqualified*
- less pollution caused by burning fuels

**or**

less specified type of pollution caused by producing heat / burning fuels

*allow correct specified pollutants caused by burning fossil fuels eg  
CO<sub>2</sub> / greenhouse gases or correct effect of burning fossil fuels eg  
global warming*

*accept thermal / heat pollution*

- using less fuel conserves resources

*accept sustainable*

*accept fossil fuels are non-renewable*


2

[7]

**Q1.** Marble chips are mainly calcium carbonate ( $\text{CaCO}_3$ ).

A student investigated the rate of reaction between marble chips and hydrochloric acid (HCl).

**Figure 1** shows the apparatus the student used.



- (a) Complete and balance the equation for the reaction between marble chips and hydrochloric acid.



(2)


- (b) The table below shows the student's results.

Time in s	Volume of gas in $\text{dm}^3$
0	0.000
30	0.030
60	0.046
90	0.052
120	0.065
150	0.070
180	0.076
210	0.079
240	0.080
270	0.080

On **Figure 2**:

- Plot these results on the grid.
- Draw a line of best fit.

**Figure 2**



(4)

- (c) Sketch a line on the grid in **Figure 2** to show the results you would expect if the experiment was repeated using 20 g of smaller marble chips.

Label this line **A**.

(2)

- (d) Explain, in terms of particles, how and why the rate of reaction changes during the reaction of calcium carbonate with hydrochloric acid.

.....


.....  
.....  
.....  
.....  
.....

(4)

(e) Another student investigated the rate of reaction by measuring the change in mass.

**Figure 3** shows the graph plotted from this student's results.

**Figure 3**



Use **Figure 3** to calculate the mean rate of the reaction up to the time the reaction is complete.

Give your answer to three significant figures.

.....

.....  
.....  
.....  
.....  
.....  
.....  
.....

Mean rate of reaction = ..... g / s

(4)

(f) Use **Figure 3** to determine the rate of reaction at 150 seconds.

Show your working on **Figure 3**.

Give your answer in standard form.

.....  
.....  
.....  
.....  
.....


Rate of reaction at 150 s = ..... g / s

(4)

(Total 20 marks)

**Q2.** A student investigated the reaction between magnesium metal and dilute hydrochloric acid.

The student placed 25 cm<sup>3</sup> of dilute hydrochloric acid in a conical flask and set up the apparatus as shown in the diagram.




The student:

- took the bung out of the flask and added a single piece of magnesium ribbon 8 cm long
- put the bung back in the flask and started a stopwatch
- recorded the volume of gas collected after 1 minute
- repeated the experiment using different temperatures of acid.

The student plotted his results on a graph.






(a) Write the correct state symbols in the equation.

Choose from (s) for solid, (l) for liquid, (g) for gas and (aq) for aqueous.



(2)

(b) The diagram shows a gas syringe after 1 minute.



(i) What volume of gas has been collected in the gas syringe after 1 minute?

Volume = ..... cm<sup>3</sup>

(1)

(ii) Use the graph to determine the temperature of the acid used in this experiment.

Temperature = ..... °C

(1)


(iii) Calculate the average rate of reaction, in  $\text{cm}^3$  of hydrogen made per second ( $\text{cm}^3/\text{s}$ ), for this experiment.

.....  
.....

Rate of reaction = .....  $\text{cm}^3/\text{s}$

(2)

(c) The student's graph has been reprinted to help you answer this question.



One of the results on the graph is anomalous.

(i) Draw a circle on the graph around the anomalous point.

(1)

(ii) Suggest what may have happened to cause this anomalous result.

Explain your answer.

.....

.....

.....

.....

.....

.....

(2)

(d) Explain how the student could improve the accuracy of the volume of gas recorded at each temperature.

.....

.....

.....

.....


.....

.....

(3)

(e) The student then used the same apparatus to measure the volume of gas produced every 10 seconds at 40 °C.

The student's results are shown on the graph.



The rate at which the gas was produced got faster over the first 60 seconds.

The student's teacher gave two possible explanations of why the reaction got faster.

**Explanation 1**

There was a layer of magnesium oxide on the surface of the magnesium. The layer of magnesium oxide prevented the magnesium reacting with the acid. As the magnesium oxide reacted slowly with the acid, the magnesium was exposed to the acid and hydrogen gas was produced.

**Explanation 2**

The reaction is exothermic, and so the temperature of the acid increased during the reaction.

- (i) Describe further experimental work the student could do to see if **Explanation 1** is correct.

.....

.....

.....

.....

.....

(2)

(ii) Describe further experimental work the student could do to see if **Explanation 2** is correct.

.....

.....

.....

.....

.....

(2)  
(Total 16 marks)

**Q3.** This question is about gold (Au).

(a) An atom of gold is represented as:



How many neutrons are in this atom of gold? .....

(1)

(b) Gold ions are used as a catalyst.

How does a gold atom (Au) become a gold ion (Au<sup>3+</sup>)?

.....  
.....  
.....  
.....

(2)

(c) A gold catalyst can be used when carbon monoxide reacts with oxygen to make carbon dioxide.

(i) Complete and balance the equation for this reaction.



(2)

(ii) Carbon dioxide has a very low boiling point.

Explain why.

.....  
.....  
.....  
.....

.....  
.....


(3)

- (d) Gold is used as a catalyst in industrial processes. Gold is rare and increasingly expensive.  
Suggest **three** reasons why gold is still used in industrial processes.

.....  
.....  
.....  
.....  
.....  
.....

(3)  
(Total 11 marks)

**Q4.** The equation for a reaction to produce hydrogen is:



(a) Explain why changing the pressure does **not** affect the yield of hydrogen at equilibrium.

.....  
.....

(1)

(b) Suggest why the best yield of hydrogen at equilibrium is obtained at **low** temperatures.

.....  
.....

(1)

(c) The temperature used in industry needs to be high enough for the reaction to take place quickly. Explain, in terms of particles, why the rate of reaction increases when the temperature is increased.

.....  
.....  
.....  
.....  
.....  
.....

(3)



- (d) Scientists have developed catalysts which allow the reaction to take place quickly at lower temperatures. How could this be good for the manufacturer and for the environment?

.....

.....

.....

.....

(2)  
(Total 7 marks)

**M1.(a)** reversible

*allow equilibrium*

**1**

(b) The colour changed from blue to pink

**1**

(c) 8.3 (°C)

**1**

(d) endothermic

*allow dehydration*

*ignore reversible*

**1**

**[4]**

**M2.(a)** natural gas

*allow correct answer shown in box if answer line blank*

1

(b) (i) 450

*allow correct answer shown in box if answer line blank*

1

(ii) iron

*allow correct answer shown in box if answer line blank*

1

(iii) The catalyst lowers the activation energy.

1

(c) (the gases are) cooled

1

ammonia condenses

*allow ammonia liquefies*

1

nitrogen and hydrogen are recycled

*if no other mark awarded allow ammonia is separated for 1 mark*

1

[7]

M3. (a) reversible 1

(b) (from blue) to pink  
*do not accept*  
*incorrect initial colour* 1

(c) sensible answers such as:

- stop water reaching papers  
*accept stop entry of moisture / wet / dampness / condensation*
- water (vapour) in air  
*ignore references to toxicity of cobalt chloride*

1

[3]

<b>M4.</b>	(a)	(i)	10	1
		(ii)	OH <sup>-</sup>	1
	(b)	(i)	air	1
		(ii)	particles move faster	1
			particles collide more often	1
		(iii)	catalyst(s)	1
	(c)		liquid	1

[7]

M5.	(a)	22	1
	(b)	(i) exothermic	1
		(ii) C	1
		gives out most heat energy <i>accept has largest temperature change / increase</i> <i>allow has highest (final) temperature <b>or</b> hottest</i>	1
	(c)	(i) increases	1
		(ii) blue <i>ignore pale / dark etc</i>	1
		(iii) reversible (reaction) <i>allow goes both ways <b>or</b> two / either way</i>	1
		(iv) <u>anhydrous</u> copper sulfate	1

[8]

<b>M6.</b>	(a)	gases	1
		white	1
		solid	1
		ammonium chloride	1
	(b)	reversible	
		<i>allow phonetic spelling</i>	
		<i>allow goes both / two / either way(s)</i>	1

[5]

- M7.** (a) increases 1
- (b) the reaction is reversible 1
- (c) A liquid 1
- (d) recycled / reused (owtte)  
*accept returned to pump / start* 1
- [4]

- M8.** (a) white to blue  
*accept colourless to blue* 1
- (b) reversible 1
- [2]



- M9.** (a) water  
*accept H<sub>2</sub>O or 5H<sub>2</sub>O*  
 2 must be below halfway 1
- (b) the cold water / ice / cubes (owtte)  
*accept 'cooled down' or references to cold* 1
- (c) reversible reaction 1
- (d) (i) 0.87g 1
- (ii) the student made errors in weighing during  
 the experiments 1
- the student did not heat the copper sulfate for long enough in one of the experiments 1
- (e) white 1
- blue  
*allow 1 mark for blue to white* 1

**[8]**

- M10.**
- (a) (i) nitrogen + hydrogen → ammonia  
*accept full correct balanced equation* 1
- (ii) reversible (reaction) (owtte)  
*do not allow just 'backwards' (unqualified)* 1
- (iii) catalyst / speed up reaction  
*accept to lower activation energy* 1
- (iv) boiling point 1
- (v) recycled (owtte) 1
- (b) (i) used to make explosives (owtte) used to make medicines (owtte) 1
- (ii) used to make fertilisers (owtte) 1
- (c) (i) sensible answers such as  
provides workers (owtte)  
good transport links  
*ignore reference to raw materials* 1
- (ii) sensible idea 1
- linked reason  
idea  
*linked reason*  
eg escape of chemicals /fumes /waste gases / pollution  
*harmful to health / environmental damage owtte*  
*do not allow harmful / damage / smell (unqualified)*

risk of explosion

*because of high pressures / may endanger local people / dangerous*

risk of fire

*because of high temperatures / may endanger local people*

noise

*any detrimental effect on quality of life **or** night and day*

lorries / traffic

*danger / noise / pollution etc*

unsightly

*detrimental effect on quality of life / house prices / reduced tourism*


uses a lot of land

*loss of habitats*

1

[10]

**Q1.**The word equation shows the reaction between anhydrous cobalt chloride and water.



(a) Name the type of reaction shown by the sign  $\rightleftharpoons$

.....

(1)

(b) When the student added water to anhydrous cobalt chloride what happened?

.....

(1)

(c) A student measured the temperature rise when anhydrous cobalt chloride was added to water.

The student's results are shown in the table below.

	<b>Trial 1</b>	<b>Trial 2</b>	<b>Trial 3</b>
<b>Temperature rise in °C</b>	8.5	8.2	8.2

Calculate the mean temperature rise.

.....

Temperature = ..... °C

(1)

(d) When water was added to anhydrous cobalt chloride an exothermic reaction took place.

Name the type of reaction when hydrated cobalt chloride reacts to form anhydrous cobalt chloride and water.


.....

.....

(1)  
(Total 4 marks)

**Q2.**A flow diagram of the Haber process is shown below.

The Haber process produces ammonia from nitrogen and hydrogen.



(a) Use the correct answer from the box to complete the sentence.

air	limestone	natural gas
-----	-----------	-------------

Hydrogen is obtained from .....

(1)

(b) In the reactor, nitrogen and hydrogen at a high pressure are heated and passed over a catalyst.

(i) Use the correct answer from the box to complete the sentence.

25	100	450
----	-----	-----

The temperature in the reactor is ..... °C

(1)

(ii) Use the correct answer from the box to complete the sentence.

copper	iron	nickel
--------	------	--------

The catalyst used in the reactor is .....

(1)

(iii) How does a catalyst speed up a reaction?

Tick (✓) **one** box.

The catalyst lowers the activation energy.

The catalyst gives the reactants extra energy.

The catalyst increases the pressure in the reactor.

(1)

(c) A mixture of gases leaves the reactor.

The mixture contains ammonia, nitrogen and hydrogen.

Describe what happens to this mixture of gases in the condenser.

Use the flow diagram to help you.

.....

.....

.....

.....


.....

.....

(3)

(Total 7 marks)

**Q3.** Read the information and then answer the questions.




**COBALT CHLORIDE PAPER**

Cobalt chloride paper can be used to test for water.

The paper contains anhydrous cobalt chloride.

The jar containing the papers must be kept closed when not being used.

The equation shows the reaction between anhydrous cobalt chloride and water.



(a) Choose **one** word from the box to complete the sentence.

<b>endothermic</b>	<b>exothermic</b>	<b>reversible</b>
--------------------	-------------------	-------------------

The symbol  $\rightleftharpoons$  means that the reaction is .....

(1)

(b) Describe the colour change when water is added to the cobalt chloride paper.

.....  
.....

(1)



(c) Suggest why the jar containing the unused cobalt chloride papers must be kept closed.

.....  
.....

(1)  
(Total 3 marks)

**Q4.** (a) Ammonia solution is used in cleaning products to remove grease from kitchen surfaces.



Ammonia solution is alkaline.

(i) Draw a ring around the number most likely to be the pH of ammonia solution.

**1**      **3**      **7**      **10**


**(1)**

(ii) Draw a ring around the ion in ammonia solution which makes it alkaline.

**Cl<sup>-</sup>**      **H<sup>+</sup>**      **Na<sup>+</sup>**      **OH<sup>-</sup>**

**(1)**

(b) Ammonia is made using the Haber process.



(i) Where does the nitrogen used in the Haber process come from?

Draw a ring around your answer.

**air**                      **natural gas**                      **water**

(1)

(ii) A high temperature of 450 °C is used in the reactor.

Tick (✓) **two** reasons in the table which explain why high temperatures make reactions faster.

Reasons	Tick (✓)
Particles move faster	
Particles are closer together	
Particles collide more often	
Particles have less energy	

(2)

(iii) The iron in the reactor speeds up the reaction but is not used up.

What is the name given to substances that speed up the chemical reaction but which are not used up during the reaction?

.....

(1)

(c) Complete the sentence.

The condenser separates the ammonia from the unreacted nitrogen and hydrogen by turning the ammonia into a .....

(1)

(Total 7 marks)

Q5. Hand warmers use chemical reactions.



(a) The table shows temperature changes for chemical reactions **A**, **B** and **C**.

Reaction	Starting temperature in °C	Final temperature in °C	Change in temperature in °C
<b>A</b>	18	25	+ 7
<b>B</b>	17	.....	+ 5
<b>C</b>	18	27	+ 9

What is the final temperature for reaction **B**? Write your answer in the table.

(1)

(b) (i) What name is given to reactions that heat the surroundings? .....

(1)

(ii) Which reaction, **A**, **B** or **C**, would be best to use in a hand warmer?


Reaction

Give a reason why you chose this reaction.


.....  
.....

(2)

(c) A student added water to some anhydrous copper sulfate.



The equation for the reaction is shown.



The student measured the temperature before and after the reaction.

(i) The measurements showed that this reaction can be used for a hand warmer.

Draw a ring around the correct answer to complete the sentence.

When water is added to anhydrous copper sulfate the temperature

of the mixture

- |                 |
|-----------------|
| increases.      |
| decreases.      |
| stays the same. |

(1)

(ii) Anhydrous copper sulfate is white.

What colour is seen after water is added to the anhydrous copper sulfate?

.....

(1)

(iii) What does the symbol  $\rightleftharpoons$  mean?

.....

(1)

(iv) The student heated a tube containing hydrated copper sulfate.

Name the solid substance produced.

.....

(1)

(Total 8 marks)

**Q6.** Stage smoke is used for special effects at pop concerts.




By Sam Cockman [CC BY 2.0], via Flickr

Ammonium chloride can be used to make stage smoke.

Ammonium chloride is a white solid.

When heated, ammonium chloride produces white smoke which can be blown onto the stage.

The equation shows what happens when ammonium chloride is heated and cooled.



(a) The sentences explain how the smoke is made.

Draw a ring around the correct answer in each box to complete each sentence.

Use the information and the equation to help you.

When heated, ammonium chloride makes two colourless

solids.

liquids.

gases.

These are blown into the air where they cool and make a

colourless

black

solid.

liquid.



white	gas.
-------	------

which is

ammonia. ammonium chloride. hydrogen chloride.
--

(4)


(b) Complete the sentence.

The symbol  $\rightleftharpoons$  means that the reaction is .....

(1)

(Total 5 marks)

Q7. The flow diagram shows how ammonia is made.



(a) What effect, if any, does the **pump** have on the pressure of the nitrogen and hydrogen?

Draw a ring around the correct answer to complete the sentence.

The pump 

decreases
has no effect on
increases

 the pressure.

(1)

(b) The word equation for making ammonia is:



In the **reactor** only a small amount of the nitrogen and hydrogen is changed into ammonia.

Tick (✓) the reason why.

Reason why	Tick (✓)
Ammonia is formed from two elements.	
Nitrogen and hydrogen are gases.	
The reaction is reversible.	

(1)

(c) In the **cooler** the mixture of gases is cooled.

Draw a ring around the correct answer to complete the sentence.

The cooler turns the ammonia into

- a liquid.
- a solid.
- an element.

(1)



(d) What happens to the unreacted nitrogen and hydrogen from the **reactor**?

.....

.....

(1)  
(Total 4 marks)

**Q8.** The diagram shows how anhydrous copper sulfate can be used to test for water.



(a) What colour change will you see when water is added to the  $\text{CuSO}_4$ ?

Colour changes from ..... to .....

(1)

(b) Draw a ring around the meaning of the symbol  $\rightleftharpoons$

endothermic


exothermic

reversible


(1)

(Total 2 marks)

**Q9.** A student heated some hydrated copper sulfate crystals.  
The equation for this reaction is shown below.



The diagram shows the apparatus used.



(a) Name liquid **A** .....

(1)

(b) What helped the vapour to condense into liquid **A**?

.....  
 .....

(1)

(c) Put a tick (✓) next to the correct meaning of the symbol  $\rightleftharpoons$

Meaning	(✓)
equal amounts of reactants and products	
exothermic reaction	
reversible reaction	

(1)

- (d) The student weighed the copper sulfate before and after it was heated.  
The experiment was repeated and the two sets of results are shown in the table.

Mass of copper sulfate before heating in grams	Mass of copper sulfate after heating in grams	Mass lost in grams
2.50	1.65	0.85
2.50	1.61	0.89

- (i) Draw a ring around the **average** mass lost for these two sets of results.

**0.85 g      0.87 g      0.89 g**

(1)

- (ii) The student used the same mass of copper sulfate each time but the mass lost was different.

Put a tick (✓) next to the **two** reasons which could explain why the mass lost is different.

Reason	(✓)
The student used different test tubes for the two experiments.	
The student made errors in weighing during the experiments.	
The student used more ice in one of the experiments.	
The student did not heat the copper sulfate for long enough in one of the experiments.	

(2)

(e) Anhydrous copper sulfate is used to test for water.

Use words from the box to complete the sentence.

blue	green	red	white
------	-------	-----	-------


Water changes the colour of anhydrous copper sulfate from .....

to .....

(2)

(Total 8 marks)

**Q10.** The Haber process is named after the German chemist, Fritz Haber. The diagram shows the main stages in the Haber process.



*Reproduced with the permission of Nelson Thornes Ltd from PATRICK FULLICK et al, ISBN 0-7487-9644-4. First published in 2006*

(a) Use the diagram to help you to answer these questions.

(i) Complete the word equation for the reaction that takes place in the reactor.



(1)

(ii) What does the symbol  $\rightleftharpoons$  mean?

.....

(1)

(iii) What is the purpose of the iron in the reactor?

.....

(1)

(iv) Ammonia is separated from unreacted nitrogen and hydrogen.

Draw a ring around the physical property that allows this separation to take place.



**boiling point      density      melting point**

**(1)**

(v) What is done with the unreacted nitrogen and hydrogen?

.....

**(1)**

(b) Some of the products that can be made from ammonia are:

- fertilisers
- dyes
- explosives
- medicines
- plastics

(i) The Haber process was invented a few years before the start of the First World War. It is thought that the First World War would have finished earlier if the Germans had **not** invented the Haber process.

Suggest why.

.....

.....

**(1)**

(ii) The Haber process has helped to increase food production.

Explain why.

.....

.....

**(1)**

(c) Factories that make ammonia are very large and operate night and day.

(i) Ammonia factories are often near towns.

Suggest why.

.....

(1)

- (ii) Suggest and explain **one** reason why local people might not want an ammonia factory near their town.

.....

.....

.....

.....

(2)

(Total 10 marks)


- M1.** (a) to speed up the reaction **or** it is a catalyst  
*allow higher level answers such as to reduce the activation energy*  
*ignore cost or yield* 1
- (b) (i) reaction is exothermic  
*accept reverse reaction is endothermic **or** high temperature causes decomposition of ammonia*  
*ignore reference to rate* 1
- (ii) more (gaseous) reactant molecules than (gaseous) product molecules  
*accept 4 volumes / moles of reactant and 2 volumes / moles of product*  
*accept lower volume of products **or** volume lower on right hand side*  
*accept 'favours the reaction which produces fewer molecules'*  
*ignore incorrect number of moles*  
*ignore reference to 'amount' of product / reactant*  
*ignore references to rate* 1
- (c) (rate is) too slow / slower owtte  
*allow catalyst would not work*  
*accept at higher temperature the rate is quicker*  
*accept at lower temperatures particles do not collide as often **or** fewer particles have the activation energy **or** particles do not have the activation energy*  
*ignore reaction would not work*  
*ignore optimum / compromise type answers* 1
- (d) cooled  
*allow ammonia / it is turned into a liquid **or** is condensed*  
*ignore references to boiling point* 1

[5]

M2. (i) reversible (reaction) 1

(ii) (yield of ammonia) increases 1

(iii)



1

[3]

M3. (i) A = air  
B = natural gas  
*for 1 mark each* 2

(ii) nitrogen  
*both for 1 mark* 1

(iii) catalyst / speed up reaction  
*for 1 mark* 1

(iv) recycle unreacted gases / save money  
*for 1 mark* 1

[5]

- M4.** (a) as a catalyst  
*accept to speed up the reaction (equilibrium)* 1
- (b) nitrogen + hydrogen  $\rightleftharpoons$  ammonia  
 $N_2 + H_2 \rightleftharpoons NH_3$   
*accept mixed formula / word equations*  
*ignore balancing* 1
- (c) (i) the reaction is reversible / an equilibrium  
*accept that ammonia can break down*  
*again into nitrogen and hydrogen*  
*accept reaction goes both ways*  
*do **not** accept some nitrogen and*  
*hydrogen do not react* 1
- (ii) (the gases are cooled)  
*no marks as given in the diagram*  
*accept correct formulae  $NH_3$ ,  $N_2$   $H_2$*  1
- ammonia removed as a liquid  
*accept ammonia liquefies **or** condenses*
- nitrogen and hydrogen are recycled  
*accept nitrogen and hydrogen are put*  
*back through the converter*  
*accept 'other gases' only if ammonia*  
*identified for first mark* 1

[5]

- M5.** (a) endothermic (reaction)  
*accept thermal decomposition* 1
- (b) gives out heat (energy)  
*accept exothermic (reaction)* 1
- turns blue  
*accept goes to hydrated copper sulphate* 1

**[3]**

- M6.** (a) (i) gas  
*accept they are all gases* 1
- (ii) reversible (reaction)  
*accept can go either way*  
*accept ammonia can be decomposed (to nitrogen and hydrogen)*  
*accept could be (an) equilibrium*  
*do not credit just 'equilibrium'* 1
- (iii) (liquid) air **or** atmosphere 1
- (iv) same number **or** amount **or** weight (of atoms) on each side (of the equation)  
*accept "sums" for each side*  
*accept same amounts of elements on each side*  
*do not credit molecules **or** compounds*  
*do not credit both sides are the same unless explained* 1
- of the same type  
*or gives a correct example 'e.g. six hydrogen atoms' (on each side)* 1
- (b) (i) nitrate **or** sulphate **or** phosphate  
*if first left blank, second may be awarded*  
*do not credit chloride*
- nitric **or** sulphuric **or** phosphoric 1
- (only if correct above, exception is for ammonium chloride followed by hydrochloric acid (1 mark))  
*as appropriate if only the formula is given this should be credited*  
*only if it is correct in every detail i.e.  $NH_4NO_3$   $HNO_3$   $(NH_4)_2SO_4$   $H_2SO_4$*   
*accept correct name with an incorrect version of the formula*  
*do not credit a correct formula with an incorrect version of the name e.g. 'nitrate/sulphite' etc* 1
- any **one** of

\* (solution) can be sprayed (on the fields **or** crops)

*accept more even distribution*

\* dissolves in soil water **or** rain (water)

*accept soaks into soil (because soaks implies water)*

\* can be taken up by (plant) roots

*do not credit can be added to water to "feed" the plants*

1

(c) (i) elements **or** different atoms are bonded or joined **or** combined **or** reacted

*do not credit just 'atoms'*

*do not credit added **or** mixed*

1

(ii) (pairs of) electrons are shared

*do not credit an electron is shared*

1

[10]



- M7.** (a) (i) ammonia and hydrogen chloride  
*both required either order*  
*accept formulae if correct in every detail* 1
- (ii) ammonium chloride /  $\text{NH}_4\text{Cl}$   
*do not credit ammonia chloride* 1
- (iii) the fumes / gases / are poisonous / toxic  
*or ammonia and hydrogen chloride are*  
*poisonous / toxic / lethal*  
*accept just ammonia is poisonous / toxic*  
*accept just hydrogen chloride is*  
*poisonous / toxic*  
*accept vapour is poisonous / toxic*  
*do not credit just fumes are dangerous*  
*or harmful* 1
- (iv) nitrogen  
*do not credit N/N<sub>2</sub>* 1
- hydrogen  
*do not credit H/H<sub>2</sub>* 1
- molecule  
*do not credit compound or mole* 1
- covalent  
*accept single / molecular* 1
- (b) (i) proton  
neutron  
electron  
*either all three correct*

*or one or two correct  
however do not credit a response  
which is repeated*

2

(ii) protons and neutrons  
*both required in either order*


1

[10]

- M8.** (a) (i) *idea that it is*  
 a reaction in which the products can themselves react to reform the original substance or a reaction that can go in either direction  
*(allow explanation in terms of the specific reaction in the question)*  
*for 1 mark* 1
- (ii) nitrogen, hydrogen and ammonia  
*(allow formulae)*  
*for 1 mark* 1
- (b) (i) high pressure/400 atm  
 low temperature/100 °C  
*for 1 mark each* 2
- (ii) higher rate of reaction  
 good rate of production  
 or *idea* that more economic (ally viable)  
*(allow catalyst more effective at higher temperature)*  
*for 1 mark each* 2
- (c) (i) *ideas that it involves*  
 use of catalyst  
*gains 1 mark*  
 but use of platinum catalyst  
*gains 2 marks* 2  
 high temperature/900 °C  
*for 1 mark* 1
- (ii)  $2\text{NO} + \text{O}_2 \rightarrow 2\text{NO}_2$

*for 1 mark each*

1



*for 1 mark each*

1

(d) (i) references to

- transport reductions
- economic savings
- saves time
- guaranteed consumer/supplier

*for 1 mark each*

2

(ii) • selection of site

- design of plant
- safe disposal of waste
- make gas emissions safe(r)
- monitoring/safety checks
- reduction of waste gas emissions
- research into more efficient processes
- research into energy savings/use of cooling water
- training of staff re: emergency procedures
- warning/evacuation procedures for the community

(or any two sensible suggestions)


*any two for 1 mark each*

2

[15]

**Q1.** Ammonium nitrate is an important chemical. The diagram shows the main stages in the manufacture of ammonium nitrate.

Study the diagram and then answer the question.



(a) What is the purpose of the iron in reactor 1?

.....  
 .....

(1)

(b) Explain why the best yield of ammonia at equilibrium is obtained:

(i) at low temperature

.....  
 .....

(1)

(ii) at high pressure.

.....  
.....

(1)

(c) The temperature used in reactor **1** is 450 °C.

Explain why a much lower temperature is **not** used.

.....  
.....

(1)

(d) A mixture of ammonia, nitrogen and hydrogen leaves reactor **1**.


In the separator, what is done to the mixture to separate the ammonia from the other gases?

.....  
.....

(1)

(Total 5 marks)

**Q2.** Transition metals are useful as catalysts. Iron is used as a catalyst in the manufacture of ammonia.



(i) What is meant by  $\rightleftharpoons$  in the chemical equation?

.....  
.....


(1)

(ii) What would be the effect on the yield of ammonia if the pressure was increased?

.....  
.....

(1)


(iii) Draw a diagram to show the arrangement of the electrons in a molecule of ammonia. The electron arrangement of each atom is shown.



(1)

(Total 3 marks)

**Q3.** The flow chart below shows the main stages in the production of ammonium nitrate.



(i) Name the **two** raw materials shown in the flow chart as **A** and **B** by choosing words from the list.

air                  coke                  limestone                  natural gas

Raw material **A** .....

Raw material **B** .....

(2)

(ii) Complete the word equation for the reaction which makes ammonia.



..... + ..... → ammonia

(1)

(iii) What is the purpose of the iron in the reactor?

.....  
.....

(1)


(iv) What is the purpose of pipe C?

.....  
.....

(1)

(Total 5 marks)

**Q4.** The diagram shows the final stages in the manufacture of ammonia.



(a) Why is iron used in the converter?

.....  
 .....

(1)

(b) Write the word equation for the reaction in the converter.

..... + .....  $\rightleftharpoons$  .....

(1)

(c) The yield of ammonia is only about 15%.

(i) Why can the yield **not** be 100%?

.....  
 .....


(1)

(ii) Describe what happens to the mixture of gases after it leaves the converter.

.....  
.....  
.....

(2)  
(Total 5 marks)

**Q5.**Hydrated copper sulphate is a blue solid. When it is heated, white solid anhydrous copper sulphate is made. This is a reversible reaction.



- (a) To make the forward reaction work, the hydrated copper sulphate must be heated all the time.

What type of reaction is this?

.....  
.....

(1)

- (b) Anhydrous copper sulphate can be used in a test for water. What **two** things will happen when water is added to anhydrous copper sulphate?

1 .....

.....

2 .....

.....

(2)

(Total 3 marks)

**Q6.** (a) In industry ammonia is produced from nitrogen and hydrogen. The equation for the reaction is:



(i) What does the symbol (g) represent?

.....

(1)

(ii) What does the symbol  $\rightleftharpoons$  represent?

.....


(1)

(iii) Nitrogen is used for the industrial production of ammonia. From what raw material does this nitrogen come?

.....

(1)

(iv) Hydrogen is used for the industrial production of ammonia. It is obtained from the reaction between methane and steam. The equation for this reaction is:



Explain how you can tell that this equation is balanced.

.....

.....

.....

.....

(2)

(b) Ammonia is used to make ammonium salts which can be used as fertilisers.

(i) Complete the names in the following sentence.

One example is ammonium ..... which is made by reacting  
ammonia with ..... acid.

(2)

(ii) All ammonium salts are soluble in water. Why is this a useful property of a fertiliser?

.....  
.....

(1)

(c) Ammonia is a covalent, chemical compound.

(i) Complete the following sentence to describe a chemical compound.

In a chemical compound, two or more .....  
.....  
.....

(1)


(ii) What is a covalent bond?

.....  
.....


(1)

(Total 10 marks)

**Q7.** (a) The equation for the reaction that takes place when ammonium chloride is heated is:



The diagram shows how a teacher demonstrated this reaction. The demonstration was carried out in a fume cupboard.



(i) Apart from the gases normally in the atmosphere, which two gases would be at **X**?

..... and .....

(1)

(ii) Name the white solid that has formed at **Y**.

.....

(1)

(iii) Why was the demonstration carried out in a fume cupboard?

.....

.....

(1)

(iv) Complete the **four** spaces in the passage.

The chemical formula of ammonia is  $\text{NH}_3$ . This shows that there is one atom of ..... and three atoms of ..... in each ..... of ammonia. These atoms are joined by bonds that are formed by sharing pairs of electrons. This type of bond is called a ..... bond.

(4)

(b) Electrons, neutrons and protons are sub-atomic particles.

(i) Complete the **three** spaces in the table.

Name of sub-atomic particle	Relative mass	Relative charge
.....	1	+1
.....	1	0
.....	$\frac{1}{1840}$	-1

(2)

(ii) Which **two** sub-atomic particles are in the nucleus of an atom?

..... and .....

(1)

(Total 10 marks)



##

Ammonia is manufactured by the Haber Process, where nitrogen and hydrogen react together as follows:



The reaction is reversible. A balance is eventually reached when ammonia is being formed at the same rate at which it is decomposing.

This point is called 'equilibrium'.

PRESSURE (ATM)	PERCENTAGE OF AMMONIA AT EQUILIBRIUM		
	100° C	300° C	500° C
25	91.7	27.4	2.9
100	96.7	52.5	10.6
400	99.4	79.7	31.9

(a) (i) What is meant by a 'reversible reaction'?

.....  
.....

(1)

(ii) Which substances are present in the mixture at equilibrium?

.....

(1)

(b) (i) Under what conditions shown in the table is the maximum yield of ammonia obtained?

.....  
.....

(2)

- (ii) The Haber Process is usually carried out at a higher temperature than that which would produce the maximum yield. Suggest why.

.....  
.....  
.....  
.....

(2)

- (c) Ammonia can be converted into nitric acid in three stages:

Stage 1 Ammonia reacts with oxygen from the air to form nitrogen monoxide and water



Stage 2 On cooling, nitrogen monoxide reacts with oxygen from the air to form nitrogen dioxide.

Stage 3 Nitrogen dioxide reacts with water to form nitric acid and nitrogen monoxide.

- (i) Describe the conditions under which the reaction in Stage 1 takes place.

.....  
.....  
.....  
.....

(3)

(ii) Balance the equation for the reaction at Stage 2.



(1)

(iii) Balance the equation for the reaction at Stage 3.



(1)

(d) The chemical plant for manufacturing ammonia is often on the same site as plants manufacturing nitric acid and fertilisers.

(i) What advantages will this have for the manufacturing company?

.....  
.....  
.....  
.....

(2)

(ii) Briefly describe **two** important ways in which it is possible to reduce the environmental impact of such plants on the surrounding area.

1 .....

.....

2 .....

.....

(2)

(Total 15 marks)

**M1.(a)** both water vapour and ethanol will condense

*allow steam for water vapour*

*allow they both become liquids*

*allow ethane condenses at a lower temperature*

*allow some of the steam hasn't reacted*

*allow it is a reversible reaction / equilibrium*

1

(b) amount will decrease

1

because the equilibrium will move to the left

1

(c) more ethanol will be produced

1

because system moves to least / fewer molecules

1

[5]

**M2.** (a) (i) nitrogen - air  
*accept atmosphere* 1

hydrogen - north sea gas / natural gas / methane / CH<sub>4</sub>  
*accept water / (crude) oil / coal / hydrocarbons / brine* 1

(ii) *allow converse throughout*  
• high temperature gives a low yield 1

• because reaction is exothermic  
*must be linked to first bullet point* 1

• but at low temperatures the rate is (too) slow  
*if no other marks awarded accept 450°C is a compromise between yield and rate*  
**or**  
*450°C gives a reasonable yield in a reasonable time for 1 mark* 1

(iii) nitric (acid)  
*accept HNO<sub>3</sub>* 1

(b) Ammonia / Haber process can be used to make fertiliser 1

with a specified economical reason

eg raw materials for Haber process readily available  
eg transport costs are lower or no need to import  
eg Haber process is a continuous process

*ignore employment / labour costs*

1

[8]

**M3.** (a) same number of (gaseous) molecules / moles / volume on both sides of the equation

*allow particles for molecules*

*do **not** accept atoms*

*ignore amount*

1

(b) (forward) reaction is exothermic

*accept reverse answer*

1

(c) any **three** from:

- particles gain energy
- particles move faster  
*allow particles collide faster / quicker*  
*ignore move more / vibrate more*
- particles collide more **or** more collisions
- more of the collisions are successful **or**  
more of the particles have the activation energy **or**  
particles collide with more force / energy

3

(d) any **two** from:

- more product (obtained in shorter time)  
*accept better yield (of product)*
- less fuel needed  
*accept less energy / heat / electricity needed*  
**or**  
lower fuel costs  
*ignore cheaper unqualified*
- less pollution caused by burning fuels

**or**

less specified type of pollution caused by producing heat / burning fuels

*allow correct specified pollutants caused by burning fossil fuels eg  
CO<sub>2</sub> / greenhouse gases or correct effect of burning fossil fuels eg  
global warming*

*accept thermal / heat pollution*

- using less fuel conserves resources

*accept sustainable*

*accept fossil fuels are non-renewable*

2

[7]



- M4.** (a) fewer product molecules than reactant molecules (owtte) **or**  
*accept forward reaction produces fewer molecules*  
*accept left hand side for reactants and right hand side for products*

3 reactant molecules and 1 product  
**or** 3 volumes of gas becomes 1 volume of gas  
*accept high pressure favours the side with fewer molecules*  
*ignore references to reaction rate*

1

- (b) any **three** from:

- low temperature gives best yield  
*accept add heat as increased temperature **or** 'less' as poor yield*  
**or** high temperature gives poor yield
- because the reaction is exothermic  
*accept reverse argument if clearly expressed*
- reaction too slow at low temperature  
**or** reaction faster at high temperature  
*accept add heat and reaction goes faster*
- temperature used gives a reasonable yield  
at a fast rate / compromise explained  
*allow get less product but it takes less time*  
*for 2 marks*

3

[4]

- M5.** (a) (i) any **one** from:
- they are positive / cations
  - they are  $H^+$
  - opposite charges attract  
*ignore atom*

1

- (ii) potassium is more reactive (or reverse)  
*assume 'it' refers to hydrogen*  
*allow potassium reacts with water*  
*allow potassium is very reactive **or** most reactive metal / element*  
*allow hydrogen gains electrons more easily / is reduced more easily*  
*accept potassium is higher up the reactivity series*

1

- (b) **6 and 2**  
*accept correct multiples and fractions*

1

- (c) (i) the reaction / it is reversible **or** a description of a reversible reaction  
*allow 'it is an equilibrium'*  
*allow reversible symbol drawn correctly*  
*allow 'the reverse / back reaction'*

1

- (ii) **lithium nitride**

assume that 'it' or if they do not specify means lithium nitride

assume lithium / lithium nitrate refers to lithium nitride

- hydrogen is bonded / held / absorbed / has formed a compound / reacted with lithium nitride

1

plus **one** of:

- does not explode / cause a fire
- is not free / less hydrogen
- is not under pressure
- does not leak
- is only released slowly

1

- compound of hydrogen with lithium nitride / product is (more) stable / less reactive / less chance of a reaction  
*accept converse for hydrogen as below*

*assume that gas / hydrogen means gas in the cylinder*

- *hydrogen (in cylinder) / gas is not bonded / held absorbed / in a compound / reacted with lithium nitride*

1

*plus one of:*

- *can explode / cause a fire*
- *is free*
- *is under pressure*
- *can leak*
- *releases quickly*

1

- (d) (i) loss of an electron **or** loses electrons  
*do not accept any ref. to oxygen*

1

- (ii) full outer shell of 8 electrons on circle  
*need not be paired*  
*can be x, dot or e*  
*do **not** accept if extra electrons added to inner shell*

1

[10]

- M6.** (a) (i) high temperature  
*accept temperature given if  $\geq 400\text{ }^{\circ}\text{C}$   
 ignore value if "high" stated, unless silly value* 1
- endothermic or reaction takes in energy  
 or  $\Delta H$  is +ve  
*independent marks* 1
- (ii) low pressure  
*or up to and including 10 atmospheres* 1
- (low pressure) favours a reaction in which  
 more molecules are formed  
*2 moles  $\rightarrow$  4 moles  
 (2 molecules  $\rightarrow$  4 molecules)  
 independent marks* 1
- (iii) nickel **and** it is a transition / transitional  
 element / metal (owtte) **or** nickel **and**  
 variable oxidation state / number or it is  
 similar to other named transition elements  
 e.g. iron 1
- (b) (i) (bonds broken =) 2005 (kJ) 1
- (bonds formed =) 2046 (kJ) 1
- energy change =  $2005 - 2046 = (-)41$   
*for correct subtraction ignore sign* 1

(ii) (exothermic)

*if in part (b)(i) answer is not 41*

*answer is consequential on endothermic or exothermic shown*

*accept correct reasoning for **incorrect** answer from (b)(i)*

energy given out forming new bonds

*do **not** accept energy needed to form new bonds*

1

greater than energy put in to break old bonds

*accept exothermic **and** more energy given out than taken in for 1 mark*

*accept negative value for energy change **or** energy in products less than energy in reactants for 1 mark*

1

[10]

**M7.** (a) **2 marks for comments related to temperature**

low / lower / lowest temperature (or 100 °C from graph)  
*ignore references to catalyst*

1

any **one** from:

- (forward) reaction exothermic  
*or reverse reaction endothermic*
- if the temperature is increased the yield of product will decrease **or**  
reaction right to left  
*high temperature favours reverse reaction or reverse argument*  
*the lower the temperature the greater the yield = 2 marks*  
*2 marks for comments related to pressure*

1

high / higher / highest pressure (or greater than 200 atm. from graph)

1

any **one** from:

- four reactant molecules but only two product molecules (owtte)  
*reverse reaction goes from 2 molecules / moles / volumes to 4*  
*molecules / moles / volumes*
- increase in pressure favours the reaction which produces  
the least number of molecules  
*decrease in pressure favours the back reaction because it produces*  
*the most molecules*

1

(b) any **three** from:

- at low temperatures the reaction is too slow
- 450 °C gives a reasonable yield at a fast rate /  
compromise between yield and rate (\*)
- 200 atm. gives a reasonable yield at a reasonable cost / safely /  
compromise between yield and cost / safety (\*)  
*(\*) or 450°C and 200 atm / these are compromise conditions for 1*  
*mark*
- catalyst works better at higher temperature

- (very) high pressures could be dangerous (owtte)  
*safety factor*
- (very) high pressures are expensive (owtte)
- (yield is not too important because) unreacted gases can be recycled

3

[7]


- M8.** (i) a reaction in which the products can  
be changed back to reactants  
*accept a reaction that can go forwards **or** backwards* 1
- under certain conditions 1
- (ii)  $M_r \text{CaCO}_3 = 100$  1
- $M_r \text{CaO} = 56$  1
- mass of CaO = 140 (tonnes) 1
- mark consequentially*

[5]




**Q1.**In industry ethanol is produced by the reaction of ethene and steam at 300°C and 60 atmospheres pressure using a catalyst.

The equation for the reaction is:



The figure below shows a flow diagram of the process.



(a) Why does the mixture from the separator contain ethanol and water?

.....  
.....

(1)

(b) The forward reaction is exothermic.

Use Le Chatelier's Principle to predict the effect of increasing temperature on the amount of ethanol produced at equilibrium.

Give a reason for your prediction.

.....  
.....  
.....  
.....

(2)

(c) Explain how increasing the pressure of the reactants will affect the amount of ethanol produced at equilibrium.

.....

.....

.....

.....

(2)  
(Total 5 marks)

**Q2.** Humberstone was a town in the desert of Northern Chile in South America. It was built for the people who worked in the nearby sodium nitrate mines.

The sodium nitrate was used as a fertiliser.

The sodium nitrate was exported by ship to countries all around the world.


Today the mines have closed and nobody lives in Humberstone.

One of the reasons for the mines closing was the invention of the Haber process.



By Sznegra (Own work) [CC-BY-SA-3.0], via Wikimedia Commons

(a) The Haber process is used to make ammonia (NH<sub>3</sub>).



The forward reaction is exothermic.

(i) Name the raw materials that are used to supply the nitrogen and hydrogen.

Nitrogen .....

Hydrogen .....

(2)

(ii) The Haber process uses a temperature of 450 °C.

Explain, as fully as you can, why a temperature of 450 °C is used rather than a much higher temperature or a much lower temperature.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

(3)

(iii) Ammonia can be converted to ammonium nitrate by adding an acid.  
Name this acid.

.....

(1)


(b) Suggest and explain why the invention of the Haber process caused the closure of the Humberstone mines in Chile.

.....  
.....  
.....  
.....  
.....

(2)

(Total 8 marks)

**Q3.** The equation for a reaction to produce hydrogen is:



(a) Explain why changing the pressure does **not** affect the yield of hydrogen at equilibrium.

.....  
.....

(1)

(b) Suggest why the best yield of hydrogen at equilibrium is obtained at **low** temperatures.

.....  
.....

(1)

(c) The temperature used in industry needs to be high enough for the reaction to take place quickly. Explain, in terms of particles, why the rate of reaction increases when the temperature is increased.

.....  
.....  
.....  
.....  
.....  
.....

(3)

- (d) Scientists have developed catalysts which allow the reaction to take place quickly at lower temperatures. How could this be good for the manufacturer and for the environment?

.....

.....

.....

.....

(2)  
(Total 7 marks)

**Q4.** Methanol is a fuel that is used in some racing cars instead of petrol.

Methanol can be made from carbon monoxide and hydrogen. The equation for this reaction is shown below.



The forward reaction is exothermic.

(a) A high pressure (between 50 and 100 atmospheres) is used in this process.

Explain why the highest equilibrium yield of methanol is obtained at high pressure.

.....  
.....

(1)

(b) The temperature used in this process is about 250 °C.

It has been stated that, 'the use of this temperature is a compromise between the equilibrium yield of product and the rate of reaction'.

Explain this statement.


.....  
.....  
.....  
.....  
.....  
.....

(3)

(Total 4 marks)

**Q5.** Read the article and then answer the questions that follow.

**Hydrogen fuel for cars?**



Hydrogen is an excellent fuel. It can be made by the electrolysis of potassium hydroxide solution.

Hydrogen gas can be stored under pressure in a cylinder but a leak of the gas could cause an explosion.

It has been found that lithium nitride can absorb and then release large volumes of hydrogen. A chemical reaction takes place between the hydrogen and the lithium nitride. The hydrogen is held in the resulting compounds by chemical bonds.

The problem is that the rate at which hydrogen is absorbed and then released from normal sized particles of lithium nitride is slow.

Recently scientists have made 'nanosized' particles of lithium nitride. These particles absorb hydrogen in the same way as normal sized lithium nitride particles. The 'nanosized' particles have the advantage that they absorb and release the hydrogen much faster when needed in the fuel cell.

It is hoped that 'nanosized' particles of lithium nitride may provide a safe method of storing hydrogen in the future.

(a) Hydrogen is produced at the negative electrode during the electrolysis of potassium hydroxide solution.

(i) Why are hydrogen ions attracted to the negative electrode?

.....

.....

.....

(1)



(ii) Potassium ions are also attracted to the negative electrode.


Explain why hydrogen gas is formed but not potassium.

.....  
.....  
.....

(1)

(b) Lithium nitride is made by reacting lithium with nitrogen.

Balance the equation for this reaction.



(1)

(c) (i) The equation for the reaction of lithium nitride with hydrogen is:



What feature of this reaction allows the hydrogen to be released?

.....  
.....

(1)

(ii) Hydrogen stored in a fuel tank filled with lithium nitride would be safer in an accident than a cylinder full of hydrogen.

Suggest and explain why.

.....  
.....  
.....  
.....

.....

(2)

(d) Lithium nitride is an ionic compound which contains lithium ions ( $\text{Li}^+$ ) and nitride ions ( $\text{N}^{3-}$ ).

(i) The formation of a lithium ion from a lithium atom is an oxidation reaction.


Explain why.

.....


.....

(1)

(ii) The diagram shows the electronic structure of a nitrogen atom.



Complete the diagram below to show the electronic structure of a nitride ion ( $\text{N}^{3-}$ ).



(1)

(Total 8 marks)

**Q6.** The reaction of methane with steam is used in industry to make hydrogen.

(a) One of the reactions in this process is represented by this equation.



The forward reaction is endothermic.

State the conditions of temperature and pressure that would give the maximum yield of hydrogen.

Explain your answers.

(i) Temperature

.....  
.....  
.....  
.....

(2)

(ii) Pressure

.....  
.....  
.....  
.....

(2)

(iii) Which one of the following metals is most likely to be a catalyst for this process?  
Draw a ring around your answer.


**aluminium      lead      magnesium      nickel      sodium**

Give a reason for your choice.

.....  
 .....

(1)

(b) A second stage in this process is represented by this equation.



(i) Use the bond energies given in the table to help you to calculate the nett energy transfer (energy change) for this reaction.

Bond	Bond energy in kJ/mol
C ≡ O	1077
C = O	805
H – H	436
O – H	464

.....  
 .....

Nett energy transfer = ..... kJ/mol

(3)

(ii) State whether this reaction is exothermic or endothermic. ....

Explain, by reference to your calculation, how you know.

.....

.....

.....


.....

(2)  
(Total 10 marks)


**Q7.** Ammonia is made from nitrogen and hydrogen in the Haber process.



**Flow Chart for the Haber Process**



Effect of temperature and pressure on the amount of ammonia at equilibrium



- (a) Use the information given above and your knowledge of the Haber process and reversible reactions to help you to answer this question.

State which conditions of temperature and pressure would give the highest percentage of ammonia at equilibrium. Explain why.

.....

.....

.....

.....

.....

.....

.....

.....

(4)

(b) The Haber process uses a temperature of 450 °C and a pressure of 200 atmospheres.

Explain why these conditions are chosen.

.....

.....

.....

.....



.....

.....

(3)  
(Total 7 marks)



**Q8.** Limestone is a useful mineral. Every day, large amounts of limestone are heated in limekilns to produce lime. Lime is used in the manufacture of iron, cement and glass and for neutralising acidic soils.



(i) The decomposition of limestone is a *reversible* reaction. Explain what this means.

.....

.....

.....

.....

(2)

(ii) Calculate the mass of lime, CaO, that would be produced from 250 tonnes of limestone, CaCO<sub>3</sub>.

Relative atomic masses: C 12; O 16; Ca 40.

.....

.....

.....

.....  
.....

Mass of lime = ..... tonnes

(3)

(Total 5 marks)